

# THE DISTRIBUTION OF GALOIS ORBITS OF POINTS OF SMALL HEIGHT IN TORIC VARIETIES

JOSÉ IGNACIO BURGOS GIL, PATRICE PHILIPPON, JUAN RIVERA-LETELIER,  
AND MARTÍN SOMBRA

ABSTRACT. We study the distribution of Galois orbits of points of small height on proper toric varieties, and its application to the Bogomolov problem.

We introduce the notion of monocritical toric metrized divisor. We prove that a toric metrized divisor  $\overline{D}$  on a proper toric variety  $X$  over a global field  $\mathbb{K}$  is monocritical if and only if for every generic  $\overline{D}$ -small sequence of algebraic points of  $X$  and every place  $v$  of  $\mathbb{K}$ , the sequence of their Galois orbits on the analytic space  $X_v^{\text{an}}$  converges to a measure. When this is the case, the limit measure is a translate of the natural measure on the compact torus sitting in the principal orbit of  $X$ .

The key ingredient is the study of the  $v$ -adic modulus distribution of Galois orbits of generic  $\overline{D}$ -small sequences of algebraic points. In particular, we characterize all their cluster measures.

We generalize the Bogomolov problem by asking when a closed subvariety of the principal orbit of a proper toric variety that has the same essential minimum than the ambient variety, must be a translate of a subtorus. We prove that the generalized Bogomolov problem has a positive answer for monocritical toric metrized divisors, and we give several examples of toric metrized divisors for which the Bogomolov problem has a negative answer.

## CONTENTS

1. Introduction	1
2. Galois orbits, height of points and essential minimum	8
3. Auxiliary results on convex analysis	14
4. Modulus distribution	22
5. Equidistribution of Galois orbits and the Bogomolov property	32
6. Examples	40
7. Potential theory on the projective line and small points	47
References	53

## 1. INTRODUCTION

The study of the limit distribution of Galois orbits of points of small height was initiated by Szpiro, Ullmo and Zhang in their seminal paper [SUZ97]. For an Abelian variety defined over a number field and over an Archimedean place, they proved the equidistribution of the Galois orbits of sequences of points whose Néron-Tate height converges to zero. This equidistribution result was motivated by

---

*Date:* January 2, 2018.

*2010 Mathematics Subject Classification.* Primary 11G50; Secondary 11G35, 14G40, 14M25, 52A41.

*Key words and phrases.* Equidistribution, Galois orbits, metrized divisor, height, essential minimum, toric variety, convex analysis, potential theory.

Burgos Gil was partially supported by the MINECO research projects MTM2013-42135-P, MTM2016-79400-P and ICMAT Severo Ochoa SEV-2015-0554. Philippon was partially supported by the CNRS research project PICS 6381 “Géométrie diophantienne et calcul formel” and the ANR research project “Hauteurs, modularité, transcendance”. Burgos Gil and Philippon were also partially supported by the FP7-MC-IRSES project n° 612534 “MODULI”. Rivera-Letelier was partially supported by FONDECYT grant 1141091 and NSF grant DMS-1700291. Sombra was partially supported by the MINECO research projects MTM2012-38122-C03-02 and MTM2015-65361-P and through the “María de Maeztu” program for units of excellence MDM-2014-0445.

the Bogomolov conjecture on Abelian varieties, and eventually led to an affirmative solution by Ullmo [Ull98] and Zhang [Zha98], see also [Cin11, Ghi09, Gub07, Yam13, Yam16] for similar results in the function field case.

This equidistribution result has been widely generalized. In particular, it has been extended to more general varieties and height functions and, with the introduction of Berkovich spaces, to non-Archimedean places [Bil97, Cha00, FR06, Cha06, BR06, Yua08, BB10, Che11]. However, all these generalizations are restricted to height functions that satisfy a special condition, namely, that the essential minimum of the heights of points is equal to the normalized height of the ambient variety, see below for precisions. In this paper, a height function satisfying this extremal condition is called “quasi-canonical”. All the available methods to prove equidistribution for points of small height break down for height functions that are not quasi-canonical.

There are important classes of quasi-canonical height functions, such as Néron-Tate heights on Abelian varieties, canonical metrics on toric varieties, and more generally those coming from algebraic dynamical systems. But there are also many height functions of interest that are not quasi-canonical, like (twisted) Fubini-Study heights on projective spaces and the Faltings height on modular varieties.

For toric varieties and height functions the situation is startling: the only ones that are quasi-canonical are essentially the canonical one, and those derived from it by scaling and translations. So all the previous equidistribution results apply to a very restricted class of toric height functions. In this paper, we give a complete description of the equidistribution phenomenon for general toric heights. Our results reveal that a very mild positivity assumption is enough to guarantee equidistribution, see Corollary 1.2 and Theorem 6.4 for restricted applications. This provides a wealth of new height functions for which the equidistribution property holds. Moreover, we give a complete classification of those toric heights for which equidistribution holds (Theorem 1.1), and use it to prove that the equidistribution property implies the Bogomolov property in the toric context (Theorem 1.4). As a by-product, we give a characterization of those toric heights whose essential minimum is attained (Corollary 4.9). We also provide examples of toric height functions that fail the Bogomolov property and for which the equidistribution property fails in a myriad of ways (§6 and §7).

Our methods build on the results and techniques developed in [BPS14, BMPS16, BPS15] to study toric heights. In particular, convex analysis and the Legendre-Fenchel duality play an important role. We introduce new techniques to deal with the spaces of adelic measures that appear naturally in the equidistribution problem. Most of the technical difficulties arise from the fact that these spaces are not compact. In dealing with these difficulties we are naturally led to consider the interplay between several topologies on these spaces.

To describe our results more precisely, we start with a brief review of the state of the art in the general setting. Let  $\mathbb{K}$  be a global field, that is, a field that is either a number field or the function field of a regular projective curve over an arbitrary field, and  $\mathfrak{M}_{\mathbb{K}}$  its set of places. We denote  $|\cdot|_v$  and  $n_v$  the absolute value on  $\mathbb{K}$  associated to a place  $v$  and its weight. Let  $X$  be a proper algebraic variety over  $\mathbb{K}$  of dimension  $n$ , and  $\overline{D} = (D, (\|\cdot\|_v)_{v \in \mathfrak{M}_{\mathbb{K}}})$  a semipositive metrized (Cartier) divisor with  $D$  big. Let

$$h_{\overline{D}}: X(\overline{\mathbb{K}}) \longrightarrow \mathbb{R}$$

be the associated height function on the set of algebraic points of  $X$ , see §2 for details. It is a generalization of the notion of height of algebraic points considered by Weil, Northcott and others.

The *essential minimum* of  $X$  with respect to  $\bar{D}$ , denoted by  $\mu_{\bar{D}}^{\text{ess}}(X)$ , is the smallest possible limit value of the height of a generic net of algebraic points of  $X$ . Consequently, we say that a net  $(p_l)_{l \in I}$  is  $\bar{D}$ -small if

$$\lim_l h_{\bar{D}}(p_l) = \mu_{\bar{D}}^{\text{ess}}(X).$$

A fundamental inequality by Zhang [Zha95] shows that the essential minimum can be bounded below in terms of the height and the degree of  $\bar{D}$ :

$$\mu_{\bar{D}}^{\text{ess}}(X) \geq \frac{h_{\bar{D}}(X)}{(n+1) \deg_D(X)}. \quad (1.1)$$

We say that  $\bar{D}$  is *quasi-canonical* if this lower bound for the essential minimum is an equality (Definition 2.7).

For a place  $v \in \mathfrak{M}_{\mathbb{K}}$ , we denote by  $X_v^{\text{an}}$  the  $v$ -adic analytification of  $X$ . If  $v$  is Archimedean, it is a complex analytic space whereas, if  $v$  is non-Archimedean, it is a Berkovich space over  $\mathbb{C}_v$ , the completion of the algebraic closure of the local field  $\mathbb{K}_v$ . We endow the space of probability measures on  $X_v^{\text{an}}$  with the weak-\* topology with respect to the space of continuous functions on  $X_v^{\text{an}}$ .

For an algebraic point  $p$  of  $X$ , we denote by  $\text{Gal}(p)_v$  its  $v$ -adic Galois orbit, that is, the orbit of  $p$  in  $X_v^{\text{an}}$  under the action of the absolute Galois group of  $\mathbb{K}$ . We set

$$\mu_{p,v} = \frac{1}{\#\text{Gal}(p)_v} \sum_{q \in \text{Gal}(p)_v} \delta_q \quad (1.2)$$

for the uniform probability measure on  $\text{Gal}(p)_v$ . We also denote by  $c_1(D, \|\cdot\|_v)^{\wedge n}$  the  $v$ -adic Monge-Ampère measure of  $\bar{D}$ , see for instance [BPS14, §1.4]. It is a measure on  $X_v^{\text{an}}$  of total mass  $\deg_D(X)$ .

The following statement is representative of several equidistribution theorems for Galois orbits of small points in the literature. In this form, it is due to Yuan [Yua08, Theorem 3.1] for number fields and to Gubler [Gub08, Theorem 1.1] for function fields, see also [Fab09] for this latter case.

**Theorem 1** (Equidistribution for quasi-canonical metrics). *Let  $X$  be a projective variety over  $\mathbb{K}$  of dimension  $n$ , and  $\bar{D}$  a quasi-canonical semipositive metrized divisor on  $X$  with  $D$  ample. Let  $(p_l)_{l \in I}$  be a generic  $\bar{D}$ -small net of algebraic points of  $X$ . Then, for every  $v \in \mathfrak{M}_{\mathbb{K}}$ , the net of probability measures  $(\mu_{p_l,v})_{l \in I}$  converges to  $\frac{1}{\deg_D(X)} c_1(D, \|\cdot\|_v)^{\wedge n}$ , the normalized  $v$ -adic Monge-Ampère measure of  $\bar{D}$ .*

A common feature of this result and its variants and generalizations, is the assumption that the lower bound (1.1) is an equality or, in other words, that the metrized divisor  $\bar{D}$  is quasi-canonical. This severely restricts their range of application. Nonetheless, these results do apply to the important case of metrics arising from algebraic dynamical systems and, moreover, they have a very strong thesis: not only the Galois orbits of points of small height do converge, but the limit measure is given by the normalized  $v$ -adic Monge-Ampère measure.

The motivation of this paper is to start the study of what happens when we remove the hypothesis that  $\bar{D}$  is quasi-canonical. Some of our typical questions are: is there always an equidistribution phenomenon for Galois orbits of  $\bar{D}$ -small points? If not, can we give conditions on  $\bar{D}$ , beyond being quasi-canonical, under which such a phenomenon occurs? When equidistribution occurs, can we describe the limit measure?

We address these questions and some of its continuations in the toric setting. As mentioned previously, our approach is based on the techniques developed in the series of papers [BPS14, BMPS16, BPS15]. We briefly recall the setting of these papers.

Let  $X$  be a proper toric variety over  $\mathbb{K}$  of dimension  $n$ , given by a complete fan  $\Sigma$  on a vector space  $N_{\mathbb{R}} \simeq \mathbb{R}^n$ , and a nef and big toric divisor  $D$  on  $X$ , given by a concave support function  $\Psi_D: N_{\mathbb{R}} \rightarrow \mathbb{R}$ . This toric divisor also defines an  $n$ -dimensional polytope  $\Delta_D$  in the dual space  $M_{\mathbb{R}} := N_{\mathbb{R}}^{\vee}$ .

Let  $\overline{D} = (D, (\|\cdot\|_v)_{v \in \mathfrak{M}_{\mathbb{K}}})$  be a semipositive toric metrized divisor on  $X$  with underlying divisor  $D$ . To it we associate an adelic family of concave functions  $\psi_{\overline{D},v}: N_{\mathbb{R}} \rightarrow \mathbb{R}$ ,  $v \in \mathfrak{M}_{\mathbb{K}}$ , called the *metric functions* of  $\overline{D}$ . They satisfy that  $|\psi_{\overline{D},v} - \Psi_D|$  is bounded on  $N_{\mathbb{R}}$  for all  $v$ , and that  $\psi_{\overline{D},v} = \Psi_D$  for all  $v$  except for a finite number. We also associate to  $\overline{D}$  an adelic family of continuous concave functions on the polytope  $\vartheta_{\overline{D},v}: \Delta_D \rightarrow \mathbb{R}$ ,  $v \in \mathfrak{M}_{\mathbb{K}}$ , called the *local roof functions* of  $\overline{D}$ . They verify that  $\vartheta_{\overline{D},v}$  is the zero function for all  $v$  except for a finite number. The *global roof function* is a concave function  $\vartheta_{\overline{D}}: \Delta_D \rightarrow \mathbb{R}$  defined as the weighted sum of the local roof functions.

Let  $\mathbb{T} \simeq \mathbb{G}_{m,\mathbb{K}}^n$  be the torus of  $X$ , which can be identified with  $X_0$ , the principal open subset of  $X$ . There is a valuation map  $\text{val}_v: \mathbb{T}_v^{\text{an}} \rightarrow N_{\mathbb{R}}$ , defined, in any given splitting of  $\mathbb{T}$ , by

$$\text{val}_v(x_1, \dots, x_n) = (-\log |x_1|_v, \dots, -\log |x_n|_v), \quad (1.3)$$

see also [BPS14, Formula (4.1.2)]. There is a canonical toric section  $s$  of  $\mathcal{O}(D)$  with  $\text{div}(s) = D$ . The metric function  $\psi_{\overline{D},v}$  is characterized by the property

$$\psi_{\overline{D},v}(\text{val}_v(p)) = \log \|s(p)\|_v$$

for  $p \in X_0^{\text{an}}$ , while the local roof function  $\vartheta_{\overline{D},v}$  is defined as the Legendre-Fenchel dual of  $\psi_{\overline{D},v}$ . We use the extension of these constructions to the case of  $\mathbb{R}$ -divisors, see §2 and [BMPS16, §4] for precisions.

The metric functions and the roof functions convey a lot of information about the pair  $(X, \overline{D})$ . For instance, the essential minimum of  $X$  with respect to  $\overline{D}$  can be computed as the maximum of the global roof function [BPS15, Theorem A]:

$$\mu_{\overline{D}}^{\text{ess}}(X) = \max_{x \in \Delta_D} \vartheta_{\overline{D}}(x). \quad (1.4)$$

In the toric setting, the condition that the metrized divisor  $\overline{D}$  is quasi-canonical is very restrictive, since it is equivalent to the condition that its global roof function is constant (Proposition 5.3). Thus, the only toric metrics to which Theorem 1 applies are those whose global roof function is constant.

To identify the toric metrics having good equidistribution properties, we introduce the notion of *monocritical* toric metrized divisor. To define this concept, first consider the map from  $X_0(\overline{\mathbb{K}})$  to the space of measures on the adelic space  $\bigoplus_{v \in \mathfrak{M}_{\mathbb{K}}} N_{\mathbb{R}}$  given by

$$p \longmapsto \nu_p = ((\text{val}_v)_* \mu_{p,v})_{v \in \mathfrak{M}_{\mathbb{K}}},$$

where  $(\text{val}_v)_* \mu_{p,v}$  denotes the direct image under the  $v$ -adic valuation map in (1.3) of the uniform probability measure on  $\text{Gal}(p)_v$  in (1.2). For a certain metric space  $\mathcal{H}_{\mathbb{K}}$  of measures defined on  $\bigoplus_{v \in \mathfrak{M}_{\mathbb{K}}} N_{\mathbb{R}}$  we show that there is a (Lipschitz) continuous function  $\eta_{\overline{D}}: \mathcal{H}_{\mathbb{K}} \rightarrow \mathbb{R}$  extending the height function  $h_{\overline{D}}$  in the sense that for every  $p$  in  $X_0(\overline{\mathbb{K}})$  we have  $h_{\overline{D}}(p) = \eta_{\overline{D}}(\nu_p)$ , see §4 for precisions. We show that this function always attains its minimum value and we give a characterization of the set of measures at which this function is minimized (Lemma 4.8 and Corollary 4.10). The semipositive toric metrized divisor  $\overline{D}$  is monocritical if the function  $\eta_{\overline{D}}$  attains its minimum at a unique measure (Definition 4.14, see also Proposition 4.16 for equivalent formulations). For such a toric metrized divisor  $\overline{D}$ ,

the uniquely minimizing measure is supported on a single point

$$\mathbf{u} = (u_v)_{v \in \mathfrak{M}_{\mathbb{K}}} \in \bigoplus_{v \in \mathfrak{M}_{\mathbb{K}}} N_{\mathbb{R}}$$

that satisfies  $\sum_v n_v u_v = 0$ , which is called the *critical point* of  $\overline{D}$  (Corollary 4.10).

The condition for  $\overline{D}$  of being monocritical can be characterized in terms of its global roof function: given a point  $x_{\max} \in \Delta_D$  maximizing  $\vartheta_{\overline{D}}$ , the sup-differential  $\partial\vartheta_{\overline{D}}(x_{\max})$  is a convex subset of  $N_{\mathbb{R}}$  containing the point 0. Then  $\overline{D}$  is monocritical if and only if 0 is a vertex of this convex subset and, when this is the case, the critical point of  $\overline{D}$  can be computed from the sup-differential of the local roof functions at  $x_{\max}$  (Proposition 4.16).

For each  $v \in \mathfrak{M}_{\mathbb{K}}$ , we denote by  $\mathbb{S}_v$  the compact subtorus of  $\mathbb{T}_v^{\text{an}}$ . To a monocritical toric metrized divisor  $\overline{D}$  with critical point  $\mathbf{u} \in \bigoplus_{v \in \mathfrak{M}_{\mathbb{K}}} N_{\mathbb{R}}$ , we associate a probability measure  $\lambda_{\mathbb{S}_v, u_v}$  on  $X_v^{\text{an}}$  (Definition 5.1). When  $v$  is Archimedean, it is the uniform measure on a translate of  $\mathbb{S}_v \simeq (S^1)^n$  whereas, when  $v$  is non-Archimedean, it is the Dirac measure at a translate of the Gauss point of  $\mathbb{T}_v^{\text{an}}$ .

The following is the main result of this paper (Theorem 5.2).

**Theorem 1.1** (Equidistribution for general toric metrics). *Let  $X$  be a proper toric variety over  $\mathbb{K}$  and  $\overline{D}$  a semipositive toric metrized divisor on  $X$  with  $D$  big. Then  $\overline{D}$  is monocritical if and only if for every place  $v \in \mathfrak{M}_{\mathbb{K}}$  and every generic  $\overline{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$ , the net of probability measures  $(\mu_{p_l, v})_{l \in I}$  on  $X_v^{\text{an}}$  converges.*

*When this is the case, the limit measure agrees with  $\lambda_{\mathbb{S}_v, u_v}$ , where  $u_v \in N_{\mathbb{R}}$  is the  $v$ -adic component of the critical point of  $\overline{D}$ .*

Quasi-canonical toric metrized divisors are monocritical, and Theorem 1.1 reduces to Theorem 1 in this case. However, quasi-canonical metrized divisors are rare even among monocritical metrized divisors, so Theorem 1.1 produces a wealth of new examples of metrized divisors satisfying the equidistribution property that were not covered by the previous results. A concrete class of such metrized divisors are those defined over a number field  $\mathbb{K}$  with positive smooth metrics at the Archimedean places and canonical metrics at the non-Archimedean ones (Theorem 6.4). Here we state a simplified version for the case when  $\mathbb{K} = \mathbb{Q}$ .

**Corollary 1.2.** *Let  $X$  be a proper toric variety over  $\mathbb{Q}$  and  $\overline{D}$  a semipositive toric metrized  $\mathbb{R}$ -divisor with  $D$  big. We assume that the  $v$ -adic metric of  $\overline{D}$  is, when  $v$  is the Archimedean place, smooth and positive and, when  $v$  is non-Archimedean, equal to the  $v$ -adic canonical metric of  $D$ . Then  $\overline{D}$  is monocritical, and for every generic  $\overline{D}$ -small sequence  $(p_l)_{l \geq 1}$  of algebraic points of  $X_0$  and every place  $v \in \mathfrak{M}_{\mathbb{Q}}$ , the sequence  $(\mu_{p_l, v})_{l \geq 1}$  on  $X_v^{\text{an}}$  converges to the probability measure  $\lambda_{\mathbb{S}_v, 0}$ .*

This corollary covers many typical examples of metrics on toric varieties such as weighted projective spaces and toric bundles, see §6.2. For instance, let  $X = \mathbb{P}_{\mathbb{Q}}^1$  and let  $\overline{D}$  be the divisor of the point at infinity equipped with the Fubini-Study metric at the Archimedean place and the canonical metric at the non-Archimedean places. Its essential minimum is

$$\mu_{\overline{D}}^{\text{ess}}(X) = \frac{\log(2)}{2},$$

and for every generic sequence of algebraic points of  $\mathbb{P}_{\mathbb{Q}}^1$  with height converging to this quantity, its  $\infty$ -adic Galois orbits converge to the Haar probability measure on  $S^1$ , the unit circle of the Riemann sphere (Example 6.5). This is an example where equidistribution does occur, but the limit measure is *not* given by the  $v$ -adic Monge-Ampère measure as in Theorem 1.

In the other extreme, classical examples of translates of subtori with the canonical metric can behave badly with respect to equidistribution. For instance, let  $X$  be the line of  $\mathbb{P}_{\mathbb{Q}}^2$  of equation  $2z_1 - z_2 = 0$  and  $\bar{D}$  the metrized divisor on  $X$  given by the restriction of the canonical metrized divisor at infinity of  $\mathbb{P}_{\mathbb{Q}}^2$ . As explained in Example 6.1, Theorem 1.1 implies that  $\bar{D}$  does not satisfy the equidistribution property in the sense of Definition 2.9.

The key new ingredient in the proof of Theorem 1.1 is the study of the modulus distribution of the  $v$ -adic Galois orbits of  $\bar{D}$ -small nets of algebraic points.

For an algebraic point  $p \in X_0(\bar{\mathbb{K}}) = \mathbb{T}(\bar{K})$ , the direct image measure

$$\nu_{p,v} := (\text{val}_v)_* \mu_{p,v}$$

is a probability measure on  $N_{\mathbb{R}}$  that gives the modulus distribution of its  $v$ -adic Galois orbit.

To each semipositive toric metrized divisor  $\bar{D}$  with  $D$  big, we associate an adelic family of nonempty subsets of  $N_{\mathbb{R}}$

$$(B_v, F_v)_{v \in \mathfrak{M}_{\mathbb{K}}}, \quad (1.5)$$

with  $B_v \subset F_v$  (Notation 4.2). We endow the space of probability measures on  $N_{\mathbb{R}}$  with the weak-\* topology with respect to the space of bounded continuous functions on  $N_{\mathbb{R}}$ . For a probability measure  $\nu$  on  $N_{\mathbb{R}}$ , we denote by  $\text{supp}(\nu) \subset N_{\mathbb{R}}$  its support and, if  $\nu$  has finite first moment, we denote by  $E[\nu]$  its expected value.

The next result characterizes the limit behavior of the modulus distribution for  $\bar{D}$ -small nets (Theorem 4.3 and Corollary 4.12).

**Theorem 1.3.** *Let  $X$  be a proper toric variety over  $\mathbb{K}$ ,  $\bar{D}$  a semipositive toric metrized divisor on  $X$  with  $D$  big, and  $v \in \mathfrak{M}_{\mathbb{K}}$ . For every  $\bar{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points in  $X_0$ , the net of probability measures  $(\nu_{p_l, v})_{l \in I}$  has at least one cluster point. Every such cluster point is a measure  $\nu_v$  with finite first moment that satisfies*

$$\text{supp}(\nu_v) \subset F_v \quad \text{and} \quad E[\nu_v] \in B_v. \quad (1.6)$$

*Conversely, for every probability measure  $\nu_v$  on  $N_{\mathbb{R}}$  that has finite first moment and satisfies (1.6), there is a  $\bar{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$  such that  $\nu_v$  is the limit of the net  $(\nu_{p_l, v})_{l \in I}$ .*

In the situation of Theorem 1.3, when  $F_v$  consist of only one point  $u_v$ , the net  $(\nu_{p_l, v})_{l \in I}$ , representing the modulus distribution of the Galois orbits of the net of small points  $(p_l)_{l \in I}$ , converges to the measure  $\delta_{u_v}$ . In this case, we say that  $\bar{D}$  satisfies the *modulus concentration property* at the place  $v$ .

One of the main ingredients in the proof of the toric equidistribution Theorem 1.1 is the characterization of monocritical metrized divisors as those for which, for every place  $v$ , the set  $F_v$  is reduced to a single point (Proposition 4.16). Equivalently, a metrized divisor is monocritical if and only if it satisfies modulus concentration at every place. This fact allows us to attach, to each monocritical divisor  $\bar{D}$ , a new metric on  $D$  that is quasi-canonical and such that the  $\bar{D}$ -small points are also small with respect to this new metric. In this way, we obtain Theorem 1.1 as a consequence of Theorem 1.3 and Theorem 1.

In the proofs of Theorem 1.3 and Proposition 4.16, a central role is played by a family of auxiliary concave functions  $(\Phi_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$  defined on the space of measures on  $N_{\mathbb{R}}$  with finite first moment. For each place  $v$ , the function  $\Phi_v$  is nonpositive and it is defined in terms of the metric at the place  $v$ , and in terms of a certain average of the metrics at all the other places. The crucial fact is that the function  $\eta_{\bar{D}}$  extending  $h_{\bar{D}}$  vanishes at an adelic measure  $(\nu_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$  if and only if, for each place  $v$ , the function  $\Phi_v$  vanishes at  $\nu_v$ . In this way we reduce the equidistribution problem to independent maximization problems at each place (Proposition 3.9 and

Theorem 4.3). The maximization problem at a given place is solved in §3. To do this, we use that for each place  $v$  the function  $\Phi_v$  is upper-semicontinuous with respect to the weak-\* topology defined above.

In the absence of modulus concentration, there is a wealth of limit measures of  $v$ -adic Galois orbits of  $\overline{D}$ -small nets of algebraic points. For instance, consider the projective line over a number field  $\mathbb{K}$  and any adelic set  $\mathbf{E} = (E_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$  of global capacity 1, whose associated equilibrium measures are compatible with the collection of sets in (1.5) (see Theorem 7.2 for the precise condition). Using Rumely's Fekete-Szegő theorem [Rum02], we show that, for all  $v$ , the equilibrium measure of  $E_v$  can be realized as the limit measure of a sequence of  $v$ -adic Galois orbits of  $\overline{D}$ -small points (Theorem 7.2).

As we already mentioned, the original motivation in [SUZ97] to search for equidistribution results of Galois orbits of small points was to prove the Bogomolov conjecture. The Bogomolov conjecture for toric varieties can be stated as follows: let  $X$  be a toric variety over  $\mathbb{K}$  and  $\overline{D}^{\text{can}}$  an ample toric divisor on  $X$  equipped with the canonical metric. Let  $V \subset X_{0, \overline{\mathbb{K}}}$  be a closed subvariety that is not a translate of a subtorus by a torsion point. Then there exists  $\varepsilon > 0$  such that the subset of algebraic points of  $V$  of canonical height bounded above by  $\varepsilon$ , is not dense in  $V$ . Equivalently, if  $V \subset X_{0, \overline{\mathbb{K}}}$  is a closed subvariety with  $\mu_{\overline{D}^{\text{can}}}^{\text{ess}}(V) = 0$ , then  $V$  is a translate of a subtorus by a torsion point. This statement is the toric counterpart of the Bogomolov conjecture for Abelian varieties proved by Ullmo and Zhang.

This conjecture was proved by Zhang [Zha95] for number fields, and later Bilu gave a different proof using his own equidistribution theorem [Bil97]. Here we extend Bilu's equidistribution theorem (Theorem 5.7) and use it to prove the following generalization of the Bogomolov conjecture for toric varieties (Theorem 5.12).

**Theorem 1.4.** *Let  $X$  be a proper toric variety over a number field  $\mathbb{K}$  and  $\overline{D}$  a monocritical toric metrized divisor on  $X$  with critical point  $\mathbf{u} = (u_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ . Let  $V$  be a closed subvariety of  $X_{0, \overline{\mathbb{K}}}$  with*

$$\mu_{\overline{D}}^{\text{ess}}(V) = \mu_{\overline{D}}^{\text{ess}}(X).$$

*Then  $V$  is a translate of a subtorus. Furthermore, if  $u_v \in \text{val}_v(\mathbb{T}(\mathbb{K})) \otimes \mathbb{Q}$  for all  $v$ , then  $V$  is the translate of a subtorus by an algebraic point  $p$  of  $X_0$  with  $h_{\overline{D}}(p) = \mu_{\overline{D}}^{\text{ess}}(X)$ .*

A closed subvariety of  $X_{0, \overline{\mathbb{K}}}$  with

$$\mu_{\overline{D}}^{\text{ess}}(V) = \mu_{\overline{D}}^{\text{ess}}(X)$$

is called a  $\overline{D}$ -special subvariety. We say that a given toric metrized divisor  $\overline{D}$  satisfies the *Bogomolov property*<sup>1</sup> if every  $\overline{D}$ -special subvariety is a translate of a subtorus (Definition 5.11). This property is intimately related with the equidistribution property. Indeed, we give an example of a metrized divisor  $\overline{D}$  on  $\mathbb{P}_{\mathbb{Q}}^2$  such that the line of equation  $z_0 + z_1 + z_2 = 0$  is  $\overline{D}$ -special (Example 6.6). This line is certainly not a translate of a subtorus, and so  $\overline{D}$  does not satisfy the Bogomolov property. This metrized divisor is a variant of the one in Example 6.1, and does not verify modulus concentration nor equidistribution for any place of  $\mathbb{Q}$ .

These results arise several interesting questions. For instance: is it possible that a given semipositive toric metrized divisor  $\overline{D}$  satisfies the equidistribution property at one place and not at another? We study this for the projective line showing that, under a natural rationality hypothesis, the equidistribution property holds at a given place if and only if it holds at every place (Proposition 7.5). However,

<sup>1</sup>not to be confused with the property (B) introduced by Bombieri and Zannier, and studied by Amoroso, David and other authors.

this conclusion is not true without this rationality hypothesis (Remark 7.7) and we have neither settled this question for the projective line in full generality, nor treated toric varieties of higher dimension.

It would also be interesting to see if the converse of Theorem 1.4 holds: Let  $X$  be a proper toric variety with  $\dim X \geq 2$ . Given a semipositive toric metrized divisor  $\overline{D}$  on  $X$ , with  $D$  big satisfying the Bogomolov property, is  $\overline{D}$  necessarily monocritical? In Proposition 6.7 we show that this is true in a very particular case. Extending this to the general case would reinforce the link between the equidistribution and the Bogomolov properties.

The results of this paper also inspire questions for general varieties and metrized divisors. For instance, from Corollary 1.2, it is plausible to conjecture that a toric divisor equipped with a positive smooth, but not necessarily toric, Archimedean metric and canonical non-Archimedean metrics, does satisfy the equidistribution property. A puzzling question is that of computing the essential minimum, with a formula generalizing (1.4) to the general, non-toric, case. Even more challenging seems the problem of generalizing the crucial notion of monocritical metrized divisor.

Several of the results presented in this introduction hold in greater generality and their thesis are stronger. We refer to the body of the paper for these versions. The structure of the paper is as follows. In §2 we give the preliminaries on Galois orbits and height of points. In §3 we introduce the upper semi-continuous concave functional  $\Phi_v$  and study its properties. In §4 we study the modulus distribution of  $v$ -adic Galois orbits of  $\overline{D}$ -small nets of points in toric varieties. In §5 we prove the toric equidistribution theorem 1.1 and its variants, together with the Bogomolov property for monocritical toric metrized divisors. In §6 we give examples illustrating a number of phenomena, including a non-monocritical toric metrized divisor not verifying the Bogomolov property. Finally, in §7 we use potential theory to study the limit measures that appear in the absence of modulus concentration.

**Acknowledgments.** We thank Qing Liu, Ricardo Menares, Joaquim Ortega-Cerdá and Tom Tucker for useful discussions and pointers to the literature. We also thank the referee for her/his detailed remarks that helped to improve the exposition of the paper.

Part of this work was done while the authors met at the Universitat de Barcelona, the Instituto de Ciencias Matemáticas (Madrid), the Institut de Mathématiques de Jussieu (Paris), the Universidad Católica de Chile (Santiago), the Tata Institute of Fundamental Research (Mumbai) and the Beijing International Center for Mathematical Research.

## 2. GALOIS ORBITS, HEIGHT OF POINTS AND ESSENTIAL MINIMUM

By a *global field*  $\mathbb{K}$  we mean a finite extension of either  $\mathbb{Q}$  or the function field of a regular projective curve over an arbitrary field, equipped with a certain set of places, denoted by  $\mathfrak{M}_{\mathbb{K}}$ . Each place  $v \in \mathfrak{M}_{\mathbb{K}}$  is a pair consisting of an absolute value  $|\cdot|_v$  on  $\mathbb{K}$  and a positive weight  $n_v \in \mathbb{Q}_{>0}$ , defined as follows.

The places of the field of rational numbers  $\mathbb{Q}$  consist of the Archimedean and the  $p$ -adic absolute values, normalized in the standard way, and with all weights equal to 1. For the function field  $K(C)$  of a regular projective curve  $C$  over a field  $k$ , the set of places is indexed by the closed points of  $C$ . For each closed point  $v_0 \in C$ , we consider the absolute value and weight given, for  $\alpha \in K(C)^\times$ , by

$$|\alpha|_{v_0} = c_k^{-\text{ord}_{v_0}(\alpha)} \quad \text{and} \quad n_{v_0} = [k(v_0) : k],$$

with  $c_k = \#k$  if the base field  $k$  is finite and  $c_k = e$  otherwise, and where  $\text{ord}_{v_0}(\alpha)$  denotes the order of  $\alpha$  in the discrete valuation ring  $\mathcal{O}_{C, v_0}$ .



Let  $\mathbb{K}_0$  denote either  $\mathbb{Q}$  or  $K(C)$ . In the general case when  $\mathbb{K}$  is a finite extension of  $\mathbb{K}_0$ , the set of places of  $\mathbb{K}$  is formed by the pairs  $v = (|\cdot|_v, n_v)$  where  $|\cdot|_v$  is an absolute value on  $\mathbb{K}$  extending an absolute value  $|\cdot|_{v_0}$  with  $v_0 \in \mathfrak{M}_{\mathbb{K}_0}$  and

$$n_v = \frac{[\mathbb{K}_v : \mathbb{K}_{0,v_0}]}{[\mathbb{K} : \mathbb{K}_0]} n_{v_0}, \quad (2.1)$$

where  $\mathbb{K}_v$  denotes the completion of  $\mathbb{K}$  with respect to  $|\cdot|_v$ , and similarly for  $\mathbb{K}_{0,v_0}$ . This set of places satisfies the following basic properties.

**Proposition 2.1.** *Let  $\mathbb{K}_0$  denote either  $\mathbb{Q}$  or  $K(C)$ , the function field of a regular projective curve  $C$  over a field  $k$ . Let  $\mathbb{K}$  be a finite extension of  $\mathbb{K}_0$  and  $\mathfrak{M}_{\mathbb{K}}$  the associated set of places as above. Then*

- (1) *for every  $v_0 \in \mathfrak{M}_{\mathbb{K}_0}$ , we have  $\sum_{v|v_0} n_v = n_{v_0}$ ;*
- (2) *for every  $\alpha \in \mathbb{K}^\times$ , we have  $\sum_{v \in \mathfrak{M}_{\mathbb{K}}} n_v \log |\alpha|_v = 0$  (product formula).*

*Proof.* These properties are classical, see for instance [AW45, Theorems 2 and 3].

In the function field case there is a subtlety, due to the fact that a given field may have different structures of global field depending on the choice of base curve.

Let  $C$  be a regular projective curve over  $k$  and  $K(C) \hookrightarrow \mathbb{K}$  a finite extension of fields. Then there is a regular projective curve  $B$  over  $k$  and a finite morphism  $\pi: B \rightarrow C$  such that  $\mathbb{K} \simeq K(B)$  and the previous extension can be identified with  $\pi^*: K(C) \hookrightarrow K(B)$ , see for instance [Liu02, Proposition 7.3.13 and Lemma 7.3.10].

We could give to  $\mathbb{K}$  the structure of global field defined directly by the curve  $B$ , but the obtained absolute values of  $\mathbb{K}$  would not be extensions of those of  $\mathbb{K}_0$ . To remedy this, we renormalize these absolute values of  $\mathbb{K}$  and, to preserve the product formula, we also change the weights.

From the valuative criterion of properness, for each closed point  $v_0 \in C$ , the absolute values of  $\mathbb{K}$  extending  $|\cdot|_{v_0}$  are in bijection with the closed points of the fiber above  $v_0$ . Moreover, since the map  $\pi$  is finite, for each closed point  $v \in \pi^{-1}(v_0)$ , the ring  $\mathcal{O}_{B,v}$  is a finite module over  $\mathcal{O}_{C,v_0}$ . It follows from [Bou85, *Chapitre 6, Proposition 2* in §8.2 and *Théorème 2* in §8.5] that the absolute value and weight corresponding to  $v$  are given, for  $\alpha \in K(B)^\times$ , by

$$|\alpha|_v = c_k^{-\frac{\text{ord}_v(\alpha)}{e_{v/v_0}}}, \quad n_v = \frac{e_{v/v_0} [k(v) : k]}{[\mathbb{K}(B) : \mathbb{K}(C)]}, \quad (2.2)$$

with  $e_{v/v_0}$  the ramification index of  $v$  over  $v_0$ . The same results in *loc. cit.* give the formula in (1).

For the product formula in (2), we obtain from (2.2) that

$$\sum_v n_v \log |\alpha|_v = -\log(c_k) \sum_v \frac{[k(v) : k] \text{ord}_v(\alpha)}{[\mathbb{K}(B) : \mathbb{K}(C)]} = \frac{-\log(c_k)}{[\mathbb{K}(B) : \mathbb{K}(C)]} \deg(\text{div}(\alpha)) = 0,$$

because the degree of a principal divisor on  $B$  is zero, which concludes the proof.  $\square$

For  $v \in \mathfrak{M}_{\mathbb{K}}$ , we choose an algebraic closure  $\mathbb{K}_v \subset \overline{\mathbb{K}}_v$  of  $\mathbb{K}_v$ . The absolute value  $|\cdot|_v$  on  $\mathbb{K}_v$  has a unique extension to  $\overline{\mathbb{K}}_v$ . We denote by  $\mathbb{C}_v$  the completion of  $\overline{\mathbb{K}}_v$  with respect to this extended absolute value. We also choose an algebraic closure  $\overline{\mathbb{K}}$  of  $\mathbb{K}$  and an embedding  $j_v: \overline{\mathbb{K}} \rightarrow \mathbb{C}_v$ .

Let  $X$  be a variety over  $\mathbb{K}$ , that is, a reduced and irreducible separated scheme of finite type over  $\mathbb{K}$ . The elements of  $X(\overline{\mathbb{K}})$  are called the *algebraic points* of  $X$ . For  $p \in X(\overline{\mathbb{K}})$ , its *Galois orbit* is  $\text{Gal}(p) := \text{Gal}(\overline{\mathbb{K}}/\mathbb{K}) \cdot p \subset X(\overline{\mathbb{K}})$ , that is, the orbit of  $p$  under the action of the absolute Galois group of  $\mathbb{K}$ .

For  $v \in \mathfrak{M}_{\mathbb{K}}$ , we denote by  $X_{\mathbb{K}_v}^{\text{an}}$  the  $v$ -adic analytifications of  $X$  over  $\mathbb{K}_v$  and by  $X_v^{\text{an}}$  the  $v$ -adic analytifications of  $X$  over  $\mathbb{C}_v$ . If  $v$  is Archimedean, they both coincide with a complex space ( $X_{\mathbb{K}_v}$  is equipped with an anti-linear involution

if  $\mathbb{K}_v \simeq \mathbb{R}$ ). If  $v$  is non-Archimedean, they are Berkovich spaces over  $\mathbb{K}_v$  and  $\mathbb{C}_v$ , respectively. These spaces are related by ([Ber90, Corollary 1.3.6])

$$X_{\mathbb{K}_v}^{\text{an}} = X_v^{\text{an}} / \text{Gal}(\overline{\mathbb{K}_v} / \mathbb{K}_v).$$

We denote by

$$\pi_v : X_v^{\text{an}} \rightarrow X_{\mathbb{K}_v}^{\text{an}} \quad (2.3)$$

the projection.

There is a map

$$X(\mathbb{C}_v) \hookrightarrow X_v^{\text{an}}.$$

Using the chosen inclusion  $j_v : \overline{\mathbb{K}} \hookrightarrow \mathbb{C}_v$ , we obtain a map  $X(\overline{\mathbb{K}}) \hookrightarrow X(\mathbb{C}_v)$  and, by composition the previous map, an inclusion

$$\iota_v : X(\overline{\mathbb{K}}) \hookrightarrow X_v^{\text{an}}.$$

The *v-adic Galois orbit* of an algebraic point  $p \in X(\overline{\mathbb{K}})$ , denoted by  $\text{Gal}(p)_v$ , is defined as the image of  $\text{Gal}(\overline{\mathbb{K}}/\mathbb{K}) \cdot p$  under  $\iota_v$ . It is a finite subset that does not depend on the choice of the inclusion  $j_v$ . We also denote by  $\mu_{p,v}$  the uniform discrete probability measure on  $X_v^{\text{an}}$  supported on  $\text{Gal}(p)_v$ , that is,

$$\mu_{p,v} = \frac{1}{\#\text{Gal}(p)_v} \sum_{q \in \text{Gal}(p)_v} \delta_q, \quad (2.4)$$

where  $\delta_q$  is the Dirac measure at the point  $q \in X_v^{\text{an}}$ . Hence, for a continuous function  $f : X_v^{\text{an}} \rightarrow \mathbb{R}$ ,

$$\int f \, d\mu_{p,v} = \frac{1}{\#\text{Gal}(p)_v} \sum_{q \in \text{Gal}(p)_v} f(q).$$

An  $\mathbb{R}$ -divisor on  $X$  is a linear combination of Cartier divisors on  $X$  with real coefficients. A *metrized*  $\mathbb{R}$ -divisor  $\overline{D}$  on  $X$  is an  $\mathbb{R}$ -divisor  $D$  on  $X$  equipped with a quasi-algebraic family of  $v$ -adic metrics  $(\|\cdot\|_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ , see [BMPS16, §3] for details. In *loc. cit.*, for each  $v \in \mathfrak{M}_{\mathbb{K}}$  the metric  $\|\cdot\|_v$  is defined over the analytic space  $X_{\mathbb{K}_v}^{\text{an}}$ . Note that this space was denoted “ $X_v^{\text{an}}$ ” in *loc. cit.* but since we will study equidistribution problems of Galois orbits of points that are defined over varying extensions of  $\mathbb{K}$  of arbitrary large degree it is more convenient to work on the space  $X_v^{\text{an}}$  instead that in the space  $X_{\mathbb{K}_v}^{\text{an}}$ . Hence we have changed the notation accordingly. With this point of view, every object on  $X_{\mathbb{K}_v}^{\text{an}}$  will be seen as an object on  $X_v^{\text{an}}$  by taking its inverse image under the projection  $\pi_v$ . For instance let  $\overline{D}$  be a metrized  $\mathbb{R}$ -divisor on  $X$  and  $s$  a rational  $\mathbb{R}$ -section of  $D$  [BMPS16, §3]. In *loc. cit.*, the  $v$ -adic metric  $\|\cdot\|_v$  is described by a continuous function  $\|s\|_v : X_{\mathbb{K}_v}^{\text{an}} \setminus |\text{div}(s)| \rightarrow \mathbb{R}_{>0}$ . In the current paper we denote by  $\|s\|_v$  the function on  $X_v^{\text{an}} \setminus |\text{div}(s)|$  given by the composition

$$\|s(p)\|_v = \|s(\pi_v(p))\|_v.$$

Clearly this function is invariant under the action of  $\text{Gal}(\overline{\mathbb{K}_v}/\mathbb{K}_v)$ .

To a metrized  $\mathbb{R}$ -divisor  $\overline{D}$  on  $X$  we can associate a height function

$$h_{\overline{D}} : X(\overline{\mathbb{K}}) \rightarrow \mathbb{R}$$

as follows.

Given  $p \in X(\overline{\mathbb{K}})$ , choose a rational  $\mathbb{R}$ -section  $s$  of  $D$  such that  $p \notin |\text{div}(s)|$ . Choose a finite extension  $\mathbb{F}$  of  $\mathbb{K}$  such that  $p \in X(\mathbb{F})$ . For each  $w \in \mathfrak{M}_{\mathbb{F}}$  over a place  $v \in \mathfrak{M}_{\mathbb{K}}$ , we can choose an embedding  $\sigma_w : \mathbb{F} \hookrightarrow \mathbb{C}_v$  such that the restriction of the absolute value  $|\cdot|_v$  of  $\mathbb{C}_v$  agrees with  $|\cdot|_w$ . We denote also by  $\sigma_w$  the induced map  $X(\mathbb{F}) \rightarrow X_v^{\text{an}}$ .

**Definition 2.2.** Let  $X$  be a variety over  $\mathbb{K}$ ,  $\bar{D}$  a metrized  $\mathbb{R}$ -divisor on  $X$ , and  $p \in X(\bar{\mathbb{K}})$ . With the above notation, the *height* of  $p$  with respect to  $\bar{D}$  is defined as

$$h_{\bar{D}}(p) = - \sum_{w \in \mathfrak{M}_{\mathbb{F}}} n_w \log \|s \circ \sigma_w(p)\|_v.$$

The height is independent of the choice of the rational  $\mathbb{R}$ -section  $s$ , the extension  $\mathbb{F}$  and the embeddings  $\sigma_w$ .

Instead of choosing a finite extension where the point  $p$  is defined, it is possible to express the height of an algebraic point in terms of its Galois orbit.

**Proposition 2.3.** *With the previous hypothesis and notation, the height of  $p$  with respect to  $\bar{D}$  is given by*

$$h_{\bar{D}}(p) = - \sum_{v \in \mathfrak{M}_{\mathbb{K}}} \frac{n_v}{\#\text{Gal}(p)_v} \sum_{q \in \text{Gal}(p)_v} \log \|s(q)\|_v.$$

*Proof.* Choose a finite normal extension  $\mathbb{F} \subset \bar{\mathbb{K}}$  of  $\mathbb{K}$  such that  $p \in X(\mathbb{F})$ . For each  $v \in \mathfrak{M}_{\mathbb{K}}$  we denote  $\mathfrak{M}_{\mathbb{F},v}$  the set of places of  $\mathbb{F}$  above  $v$ .

Write  $G = \text{Gal}(\mathbb{F}, \mathbb{K})$  and let  $\mathbb{F}^G$  be the fixed field. Then  $\mathbb{F}/\mathbb{F}^G$  is a Galois extension with Galois group  $G$  and  $\mathbb{F}^G/\mathbb{K}$  is purely inseparable. Hence, for  $v \in \mathfrak{M}_{\mathbb{K}}$ ,

$$\frac{[\mathbb{F}_w : \mathbb{K}_v]}{[\mathbb{F} : \mathbb{K}]} = \frac{[\mathbb{F}_w : (\mathbb{F}^G)_v]}{[\mathbb{F} : \mathbb{F}^G]} = \frac{1}{\#\mathfrak{M}_{\mathbb{F},v}}.$$

Then, from the definition of the height of  $p$  in Definition 2.2 and Proposition 2.1(1), it follows that

$$\begin{aligned} h_{\bar{D}}(p) &= - \sum_{v \in \mathfrak{M}_{\mathbb{K}}} n_v \sum_{w|v} \frac{[\mathbb{F}_w : \mathbb{K}_v]}{[\mathbb{F} : \mathbb{K}]} \log \|s \circ \sigma_w(p)\|_v \\ &= - \sum_{v \in \mathfrak{M}_{\mathbb{K}}} \frac{n_v}{\#\mathfrak{M}_{\mathbb{F},v}} \sum_{w|v} \log \|s \circ \sigma_w(p)\|_v. \end{aligned} \quad (2.5)$$

The group  $G$  acts on  $\mathfrak{M}_{\mathbb{F},v}$  and, since  $p$  is defined over  $\mathbb{F}$ , also on  $\text{Gal}(p)_v$ . Both actions are transitive. Therefore, choosing  $w_0 \in \mathfrak{M}_{\mathbb{F},v}$ ,

$$\begin{aligned} \frac{1}{\#\mathfrak{M}_{\mathbb{F},v}} \sum_{w|v} \log \|s \circ \sigma_w(p)\|_v &= \frac{1}{\#G} \sum_{g \in G} \log \|s \circ \sigma_{w_0}(g(p))\|_v \\ &= \frac{1}{\#\text{Gal}(p)_v} \sum_{q \in \text{Gal}(p)_v} \log \|s(q)\|_v. \end{aligned}$$

The statement follows from this together with (2.5).  $\square$

The *essential minimum* of  $X$  with respect to  $\bar{D}$  is defined as

$$\mu_{\bar{D}}^{\text{ess}}(X) = \sup_{\substack{Y \subseteq X \\ Y \text{ closed}}} \inf_{p \in (X \setminus Y)(\bar{\mathbb{K}})} h_{\bar{D}}(p). \quad (2.6)$$

Roughly speaking, the essential minimum is the generic infimum of the height function.

**Definition 2.4.** Let  $X$  be a variety over  $\mathbb{K}$  and  $\bar{D}$  a metrized  $\mathbb{R}$ -divisor on  $X$ . A net  $(p_l)_{l \in I}$  of algebraic points of  $X$  is  $\bar{D}$ -*small* if

$$\lim_l h_{\bar{D}}(p_l) = \mu_{\bar{D}}^{\text{ess}}(X).$$

The net  $(p_l)_{l \in I}$  is *generic* if, for every closed subset  $Y \subsetneq X$ , there is  $l_0 \in I$  such that  $p_l \notin Y(\bar{\mathbb{K}})$  for  $l \geq l_0$ .

**Proposition 2.5.** *Given a variety  $X$  over  $\mathbb{K}$  and  $\overline{D}$  a metrized  $\mathbb{R}$ -divisor on  $X$ , there exists a generic  $\overline{D}$ -small net of algebraic points of  $X$ . Moreover, every generic net  $(p_i)_{i \geq 1}$  of algebraic points of  $X$  satisfies*

$$\liminf_i h_{\overline{D}}(p_i) \geq \mu_{\overline{D}}^{\text{ess}}(X).$$

*Proof.* The second statement is clear from the definition of the essential minimum.

For the first statement, let  $I$  be the set of closed subvarieties of  $X$  of pure codimension 1, ordered by inclusion. This is a directed set. For each  $Y \in I$ , denote by  $c(Y)$  its number of irreducible components and choose a point  $p_Y \in (X \setminus Y)(\overline{\mathbb{K}})$  with

$$h_{\overline{D}}(p_Y) \leq \mu_{\overline{D}}^{\text{ess}}(X) + \frac{1}{c(Y)}.$$

Clearly, the net  $(p_Y)_{Y \in I}$  is generic and  $\overline{D}$ -small.  $\square$

**Remark 2.6.** When  $\mathbb{K}$  is a number field, the collection of subvarieties of  $X$  is countable. This fact implies that a generic  $\overline{D}$ -small net contains generic  $\overline{D}$ -small sequences (although these sequences need not be subnets). Thus, Proposition 2.5 implies the existence of generic  $\overline{D}$ -small *sequences* of algebraic points in this case.

Suppose now that the variety  $X$  is proper over  $\mathbb{K}$  and of dimension  $n$ . A metrized  $\mathbb{R}$ -divisor  $\overline{D}$  on  $X$  is *semipositive* if it can be written as

$$\overline{D} = \sum_{i=1}^r \alpha_i \overline{D}_i$$

with  $\overline{D}_i$  a semipositive metrized divisor and  $\alpha_i \in \mathbb{R}_{\geq 0}$ ,  $i = 1, \dots, r$ . Recall that  $\overline{D}_i$  is semipositive if each of its  $v$ -adic metrics is a uniform limit of a sequence of semipositive smooth (respectively, algebraic) metrics in the Archimedean (respectively, non-Archimedean) case.

Given a semipositive metrized  $\mathbb{R}$ -divisor  $\overline{D}$ , we can extend the notion of height of points to subvarieties of higher dimension. In particular, the height of  $X$ , denoted by  $h_{\overline{D}}(X)$ , is defined. Moreover, for each  $v \in \mathfrak{M}_{\mathbb{K}}$  we can consider the associated  $v$ -adic Monge-Ampère measure, denoted by  $c_1(D, \|\cdot\|_v)^{\wedge n}$ . It is a measure on  $X_v^{\text{an}}$  of total mass  $\deg_D(X)$ , see for instance [BPS14, §1.4] for the case when  $D$  is a divisor. The  $v$ -adic Monge-Ampère measure of an  $\mathbb{R}$ -divisor is defined from that of divisors by polarization and multilinearity.

A theorem of Zhang shows that, when  $\mathbb{K}$  is a number field,  $D$  is an ample divisor and  $\overline{D}$  is semipositive, the essential minimum can be bounded below in terms of the height of  $X$  and the degree of  $D$  [Zha95, Theorem 5.2]:

$$\mu_{\overline{D}}^{\text{ess}}(X) \geq \frac{h_{\overline{D}}(X)}{(n+1) \deg_D(X)}. \quad (2.7)$$

This inequality can be generalized to global fields and semiample big divisors, see for instance [Gub08, Proposition 5.10].

In some cases, the inequality (2.7) is an equality. For instance, this happens for the canonical metric on divisors of toric and Abelian varieties, and for the canonical metrics coming from dynamical systems. This motivates the following definition.

**Definition 2.7.** Let  $X$  be a proper variety over  $\mathbb{K}$  of dimension  $n$ , and  $\overline{D}$  a semipositive metrized  $\mathbb{R}$ -divisor on  $X$  with  $D$  big. Then  $\overline{D}$  is *quasi-canonical* if

$$\mu_{\overline{D}}^{\text{ess}}(X) = \frac{h_{\overline{D}}(X)}{(n+1) \deg_D(X)}.$$

In other words, quasi-canonical metrized  $\mathbb{R}$ -divisors are those for which Zhang's lower bound for the essential minimum is attained.

As we will see in §5, the condition for a toric metric of being quasi-canonical is very restrictive. The following observation is a direct consequence of Proposition 2.5 and of the inequality (2.7).

**Proposition 2.8.** *Let  $X$  be a proper variety over  $\mathbb{K}$  of dimension  $n$  and  $\bar{D}$  a semipositive metrized divisor on  $X$  with  $D$  big and semiample. Then there exists a generic net  $(p_l)_{l \in I}$  of algebraic points of  $X$  with*

$$\lim_l h_{\bar{D}}(p_l) = \frac{h_{\bar{D}}(X)}{(n+1) \deg_D(X)} \quad (2.8)$$

*if and only if  $\bar{D}$  is quasi-canonical.*

We discuss now the equidistribution of Galois orbits of points of small height.

Let  $X$  be a proper variety over  $\mathbb{K}$  and  $v \in \mathfrak{M}_{\mathbb{K}}$ . We endow the space of probability measures on  $X_v^{\text{an}}$  with the weak- $*$  topology with respect to the space of continuous functions on  $X_v^{\text{an}}$ . In particular, a net of probability measures  $(\mu_l)_{l \in I}$  converges to a probability measure  $\mu$  if, for every continuous function  $f: X_v^{\text{an}} \rightarrow \mathbb{R}$ ,

$$\lim_l \int f \, d\mu_l = \int f \, d\mu.$$

**Definition 2.9.** Let  $\bar{D}$  be a metrized  $\mathbb{R}$ -divisor on  $X$ . A probability measure  $\mu$  on  $X_v^{\text{an}}$  is a *v-adic limit measure* for  $\bar{D}$  if there exists a generic  $\bar{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X$  such that the net of probability measures  $(\mu_{p_l, v})_{l \in I}$  converges to  $\mu$ . We say that  $\bar{D}$  satisfies the *v-adic equidistribution property* if, for every generic  $\bar{D}$ -small net  $(p_l)_{l \in I}$  as above, the net of measures  $(\mu_{p_l, v})_{l \in I}$  converges.

Clearly, when the *v-adic equidistribution property* holds, there exists a unique limit measure.

**Remark 2.10.** When  $\mathbb{K}$  is a number field, the analytic space  $X_v^{\text{an}}$  is metrizable and so is the space of probability measures on it. In particular, it has a nested countable basis of neighborhoods at each point. If all the generic  $\bar{D}$ -small sequences contained in a generic  $\bar{D}$ -small net converge, they must converge to the same limit. Then, using the above fact, we may strengthen Remark 2.6 showing that a generic  $\bar{D}$ -small net not converging to a given point contains a generic  $\bar{D}$ -small sequences not converging to that point. This implies that one can reduce to sequences, instead of nets, in Definition 2.9 over number fields.

In the literature there are many equidistribution theorems of Galois orbits of points of small height. All these equidistribution results deal with generic nets (or sequences when  $\mathbb{K}$  is a number field) of algebraic points satisfying the equality (2.8). In view of Proposition 2.8, the existence of such a net implies that the metric is quasi-canonical. Moreover, the condition (2.8) for this net is equivalent of being  $\bar{D}$ -small. Thus we can reformulate a general equidistribution result in the following form.

**Theorem 2.11.** *Let  $\mathbb{K}$  be a global field and  $X$  a projective variety over  $\mathbb{K}$  of dimension  $n$ . Let  $\bar{D}$  be a semipositive metrized divisor on  $X$  such that  $D$  is ample. If  $\bar{D}$  is quasi-canonical then, for every place  $v \in \mathfrak{M}_{\mathbb{K}}$ ,*

- (1)  $\bar{D}$  satisfies the *v-adic equidistribution property*;
- (2) the limit measure is the normalized Monge-Ampère measure

$$\frac{1}{\deg_D(X)} c_1(D, \|\cdot\|_v)^{\wedge n}.$$

This result is due to Yuan [Yua08, Theorem 3.1] in the number field case and, with more general hypotheses, to Gubler [Gub08, Theorem 1.1] in the function field case.

This equidistribution theorem imposes a very restrictive hypothesis, namely, that the metrized divisor  $\overline{D}$  is quasi-canonical. But it also has a very strong thesis: not only the Galois orbits of points of small height converge to a measure, but this limit measure can be identified with the normalized Monge-Ampère measure of the metrized divisor.

The main objective of this paper is to start the study of what happens when the hypothesis of  $\overline{D}$  being quasi-canonical is removed. We will work with toric varieties and toric metrics because, in this case, the tools developed previously allow us to work very explicitly. In this setting, we will see that the first statement in Theorem 2.11 holds in much great generality, but, if the metric is not quasi-canonical, the limit measure does not need to agree with the normalized Monge-Ampère measure.

### 3. AUXILIARY RESULTS ON CONVEX ANALYSIS

In this section we gather several definitions and results on convex analysis that we will use in our study of toric height functions. For a background in convex analysis, see for instance [BPS14, §2].

Let  $N_{\mathbb{R}} \simeq \mathbb{R}^n$  be a real vector space of dimension  $n$  and  $M_{\mathbb{R}} = \text{Hom}(N_{\mathbb{R}}, \mathbb{R}) = N_{\mathbb{R}}^{\vee}$  its dual. The pairing between  $x \in M_{\mathbb{R}}$  and  $u \in N_{\mathbb{R}}$  will be denoted either by  $\langle x, u \rangle$  or  $\langle u, x \rangle$ .

Following [BPS14, §2], a convex subset  $C$  is nonempty. The *relative interior* of  $C$ , denoted by  $\text{ri}(C)$ , is the interior  $C$  relative to the minimal affine subspace containing it.

Let  $C \subset M_{\mathbb{R}}$  be a convex subset and  $g: C \rightarrow \mathbb{R}$  a concave function. The *sup-differential* of  $g$  at a point  $x \in C$  is

$$\partial g(x) = \{u \in N_{\mathbb{R}} \mid \langle u, z - x \rangle \geq g(z) - g(x) \text{ for all } z \in C\}.$$

It is a closed, convex subset of  $N_{\mathbb{R}}$ , see [BPS14, §2.2]. The *stability set* of  $g$  is the convex subset of  $N_{\mathbb{R}}$  defined by

$$\text{stab}(g) = \{u \in N_{\mathbb{R}} \mid u - g \text{ is bounded below}\}.$$

The *Legendre-Fenchel dual* of  $g$  is the concave function  $g^{\vee}: \text{stab}(g) \rightarrow \mathbb{R}$  defined by

$$g^{\vee}(u) = \inf_{x \in C} \langle u, x \rangle - g(x), \quad (3.1)$$

see *ibidem*.

Let  $E \subset N_{\mathbb{R}}$  be a convex subset. A nonempty subset  $F \subset E$  is a *face* of  $E$  if every closed segment  $S \subset E$  whose relative interior has nonempty intersection with  $F$ , is contained in  $F$ .

**Lemma 3.1.** *Let  $C \subset M_{\mathbb{R}}$  be a compact convex subset and  $g_1, g_2: C \rightarrow \mathbb{R}$  two continuous concave functions. Denote by  $C_{\max}$  the convex subset of  $C$  of the points where  $g_1 + g_2$  attains its maximum value and choose  $x \in C_{\max}$ . For  $i = 1, 2$ , consider the concave function  $\widehat{\phi}_i: N_{\mathbb{R}} \rightarrow \mathbb{R}$  defined by*

$$\widehat{\phi}_i(u) = g_i^{\vee}(u) - \langle x, u \rangle + g_i(x). \quad (3.2)$$

Then

- (1) if  $x' \in \text{ri}(C_{\max})$ , then  $\partial g_i(x')$  is a face of  $\partial g_i(x)$ ,  $i = 1, 2$ ;
- (2)  $\partial g_1(x) \cap (-\partial g_2(x))$  is nonempty and does not depend on the choice of  $x \in C_{\max}$ ;
- (3) the minimal face of  $\partial g_1(x)$  containing  $\partial g_1(x) \cap (-\partial g_2(x))$  does not depend on the choice of  $x \in C_{\max}$ ;
- (4) the function  $\widehat{\phi}_i$  is nonpositive and vanishes precisely on  $\partial g_i(x)$ .

*Proof.* The restriction to  $C_{\max}$  of the sum  $g_1 + g_2$  is constant, and so the restrictions to this set of  $g_1$  and  $g_2$  are affine and with opposite slopes. In other words, there is  $u_0 \in N_{\mathbb{R}}$  such that, for all  $x_1, x_2 \in C_{\max}$ ,

$$g_1(x_2) - g_1(x_1) = \langle u_0, x_2 - x_1 \rangle \quad \text{and} \quad g_2(x_2) - g_2(x_1) = -\langle u_0, x_2 - x_1 \rangle. \quad (3.3)$$

For the statement (1), let  $i \in \{1, 2\}$ ,  $x' \in \text{ri}(C_{\max})$  and  $u \in \partial g_i(x')$ . By the definition of the sup-differential, for all  $z \in C$ ,

$$\langle u, z - x' \rangle \geq g_i(z) - g_i(x'). \quad (3.4)$$

Since  $x'$  is in the relative interior of  $C_{\max}$ , there exists  $\varepsilon > 0$  such that  $x' - \varepsilon(x - x') \in C_{\max}$ . By (3.4) and (3.3),

$$\begin{aligned} -\varepsilon \langle u, x - x' \rangle &= \langle u, x' - \varepsilon(x - x') - x' \rangle \\ &\geq g_i(x' - \varepsilon(x - x')) - g_i(x') \\ &= (-1)^{i-1} \langle u_0, -\varepsilon(x - x') \rangle = -\varepsilon(g_i(x) - g_i(x')). \end{aligned}$$

Hence  $\langle u, x - x' \rangle \leq g_i(x) - g_i(x')$ . By (3.4) applied to  $z = x$ , we have also the reverse inequality. Thus  $\langle u, x - x' \rangle = g_i(x) - g_i(x')$ , and it follows from (3.4) that, for all  $z \in C$ ,

$$\langle u, z - x \rangle \geq g_i(z) - g_i(x).$$

Hence  $u \in \partial g_i(x)$  and so  $\partial g_i(x') \subset \partial g_i(x)$ . Applying [BPS14, Proposition 2.2.8] to the closed concave function  $g_i^{\vee}$  and observing that  $g_i^{\vee\vee} = g_i$ , we deduce that  $\partial g_i(x')$  is a face of  $\partial g_i(x)$ .

To prove the statement (2) note that, since  $g_1 + g_2$  attains its maximum value at  $x$ , by [BPS14, Proposition 2.3.6(2)]

$$0 \in \partial(g_1 + g_2)(x) = \partial g_1(x) + \partial g_2(x).$$

Hence  $\partial g_1(x) \cap (-\partial g_2(x)) \neq \emptyset$ , as stated. Now let  $u$  be a point in this intersection. Then

$$\langle u, z - x \rangle \geq g_1(z) - g_1(x) \quad \text{and} \quad \langle -u, z - x \rangle \geq g_2(z) - g_2(x). \quad (3.5)$$

Choose  $x'' \in C_{\max}$ . Subtracting, from the inequalities (3.5) applied to  $z = x''$ , the identities (3.3) applied to  $x_1 = x$  and  $x_2 = x''$ , we deduce that

$$\langle u - u_0, x'' - x \rangle = 0.$$

Using this together with (3.4) and (3.5), we obtain

$$\langle u, z - x'' \rangle \geq g_1(z) - g_1(x'') \quad \text{and} \quad \langle -u, z - x'' \rangle \geq g_2(z) - g_2(x'').$$

Hence  $u \in \partial g_1(x'') \cap (-\partial g_2(x''))$ , as stated.

For the next statement, consider the convex set  $B = \partial g_1(x) \cap (-\partial g_2(x))$  that, thanks to (2), does not depend on the choice of  $x \in C_{\max}$ . Denote by  $F_x$  the minimal face of  $\partial g_1(x)$  containing it. By (1), it is enough to consider the case when  $x \in \text{ri}(C_{\max})$ . By the same statement, the set  $\partial g_2(x)$  does not depend on the choice of  $x \in \text{ri}(C_{\max})$ , proving (3).

The statement (4) follows readily from [BPS14, Lemma 2.2.6].  $\square$

**Definition 3.2.** Let  $C \subset M_{\mathbb{R}}$  be a compact convex subset and  $g_1, g_2: C \rightarrow \mathbb{R}$  two continuous concave functions. Choose a point  $x$  in  $C$  at which  $g_1 + g_2$  attains its maximum value. We define the convex subset of  $N_{\mathbb{R}}$

$$B(g_1, g_2) = \partial g_1(x) \cap (-\partial g_2(x))$$

and the convex subset

$$F(g_1, g_2) \subset \partial g_1(x)$$

as the minimal face of  $\partial g_1(x)$  that contains  $B(g_1, g_2)$ . By Lemma 3.1(2,3), these convex subsets do not depend on the choice of  $x$ .

**Lemma 3.3.** *Let  $C \subset M_{\mathbb{R}}$  be a compact convex subset with nonempty interior and  $g_1, g_2: C \rightarrow \mathbb{R}$  two concave functions. Then  $B(g_1, g_2)$  is bounded and  $F(g_1, g_2)$  contains no lines.*

*Proof.* The convex set  $B(g_1, g_2)$  is not bounded if and only if it contains a ray, that is, a subset of the form  $\mathbb{R}_{\geq 0}u_1 + u_2$  with  $u_i \in N_{\mathbb{R}}$ ,  $i = 1, 2$ , and  $u_1 \neq 0$ . Suppose that this is the case. This implies that, for  $x \in C_{\max}$  and all  $t \geq 0$ ,

$$tu_1 + u_2 \in \partial g_1(x) \quad \text{and} \quad -tu_1 - u_2 \in \partial g_2(x).$$

Hence, for all  $z \in C$  and  $t \geq 0$ ,

$$-\langle u_2, z - x \rangle + g_1(z) - g_1(x) \leq t\langle u_1, z - x \rangle \leq -\langle u_2, z - x \rangle - g_2(z) + g_2(x).$$

Letting  $t \rightarrow \infty$ , this implies  $C \subset \{z \mid \langle u_1, z - x \rangle = 0\}$ , contradicting the hypothesis that  $C$  has nonempty interior. Hence  $B(g_1, g_2)$  is bounded.

Similarly, if  $F(g_1, g_2)$  contains a line  $\mathbb{R}u_1 + u_2$ , then, for  $x \in C_{\max}$  and  $t \in \mathbb{R}$ ,

$$tu_1 + u_2 \in \partial g_1(x).$$

This also implies that  $C$  is contained in the affine hyperplane  $\{z \mid \langle u_1, z - x \rangle = 0\}$  and contradicts the hypothesis that  $C$  has nonempty interior. Hence  $F(g_1, g_2)$  contains no lines.  $\square$

Let  $\mathcal{C}_b(N_{\mathbb{R}})$  be the space of bounded continuous functions on  $N_{\mathbb{R}}$ , let  $\|\cdot\|$  be an auxiliary norm on  $N_{\mathbb{R}}$  that we fix, and for  $x$  in  $N_{\mathbb{R}}$  and  $r > 0$  denote by  $\mathcal{B}(x, r)$  the open ball in  $N_{\mathbb{R}}$  centered at  $x$  and of radius  $r$ .

**Definition 3.4.** We denote by  $\mathcal{P}$  the space of Borel probability measures on  $N_{\mathbb{R}}$  endowed with the weak-\* topology with respect to  $\mathcal{C}_b(N_{\mathbb{R}})$ . This is the coarsest topology on  $\mathcal{P}$  such that, for all  $\varphi \in \mathcal{C}_b(N_{\mathbb{R}})$ , the function  $\mu \mapsto \int \varphi \, d\mu$  is continuous.

We denote by  $\mathcal{E} \subset \mathcal{P}$  the topological subspace of probability measures with finite first moment, that is, the probability measures on  $N_{\mathbb{R}}$  satisfying

$$\int \|u\| \, d\mu(u) < \infty.$$

For  $\mu \in \mathcal{E}$ , the *expected value* is

$$E[\mu] = \int u \, d\mu(u) \in N_{\mathbb{R}}.$$

The weak-\* topology of  $\mathcal{P}$  with respect to  $\mathcal{C}_b(N_{\mathbb{R}})$  is called the “*topologie étroite*” in [Bou69, §5]. By Proposition 5.4.10 in *loc. cit.*, the topological space  $\mathcal{P}$  is complete, metrizable and separable. Later we will consider other topologies on the underlying spaces of  $\mathcal{P}$  and  $\mathcal{E}$ . When this is the case, we will state explicitly the used topology.

For  $\mu \in \mathcal{P}$ , its *support*, denoted by  $\text{supp}(\mu)$ , is the set of all points in  $N_{\mathbb{R}}$  such that all its neighborhoods have positive measure. Clearly, every measure in  $\mathcal{P}$  with bounded support lies in  $\mathcal{E}$ .

**Proposition 3.5.** *The space  $\mathcal{E}$  verifies the following properties.*

- (1) *For every  $\mu$  in  $\mathcal{E}$  we have  $E[\mu] \in \text{conv}(\text{supp}(\mu))$ .*
- (2) *The set of probability measures on  $N_{\mathbb{R}}$  with finite support is dense in  $\mathcal{E}$ .*

*Proof.* To prove the first statement, let  $\mu \in \mathcal{E}$  and suppose that  $E[\mu]$  does not lie in  $\text{conv}(\text{supp}(\mu))$ . Restricting to an affine subspace if necessary, we can assume that  $\text{conv}(\text{supp}(\mu)) \cup \{E[\mu]\}$  is not contained in a hyperplane. The hyperplane separation theorem applied to the convex sets  $\{E[\mu]\}$  and  $\text{conv}(\text{supp}(\mu))$ , implies



that there is a nonconstant affine function  $f$  such that  $f(\mathbb{E}[\mu]) \geq 0$  and  $f|_{\text{supp}(\mu)} \geq 0$ , see for example [Roc70, Theorem 11.3]. So

$$0 \geq f(\mathbb{E}[\mu]) = \int f(u) d\mu \geq 0,$$

and therefore  $\mathbb{E}[\mu]$  and  $\text{supp}(\mu)$  are both contained in the hyperplane  $\{u \in N_{\mathbb{R}} \mid f(u) = 0\}$ . This contradiction completes the proof of the first statement.

To prove the second statement, we show that every measure in  $\mathcal{E}$  is the limit of measures with bounded support. For  $r > 0$  put  $\mathcal{B}(0, r) = \{x \in N_{\mathbb{R}} \mid \|x\| \leq r\}$ . Given a measure  $\mu \in \mathcal{E}$ , the sequence of probability measures with compact support defined for  $l \geq 1$  by

$$\mu|_{\mathcal{B}(0, l)} + \mu(N_{\mathbb{R}} \setminus \mathcal{B}(0, l)) \delta_0,$$

converges to  $\mu$  as  $l \rightarrow \infty$ .

Using a straightforward discretization argument, one can show that every measure in  $\mathcal{E}$  with bounded support is the limit of probability measures with finite support. Combined with the previous observation, this completes the proof of the second statement.  $\square$

For the rest of this section, we fix a compact convex subset  $C \subset M_{\mathbb{R}}$  with nonempty interior and two continuous concave functions  $g_1, g_2: C \rightarrow \mathbb{R}$ . Since  $C$  is compact, the stability set of  $g_i$  is  $N_{\mathbb{R}}$ . Thus the Legendre-Fenchel dual  $g_i^{\vee}$  is a concave function on  $N_{\mathbb{R}}$  with stability set  $C$ .

We introduce the function  $\Phi: \mathcal{E} \rightarrow \mathbb{R}$  given, for  $\mu \in \mathcal{E}$ , by

$$\Phi(\mu) = \int g_1^{\vee} d\mu + g_2^{\vee}(-\mathbb{E}[\mu]) + \max_{x \in C} (g_1(x) + g_2(x)). \quad (3.6)$$

This function will play a central role in the proof of the main results in the next section.

It follows easily from its definition that  $\Phi$  is concave. In general, this function is not continuous, as the following example shows.

**Example 3.6.** Let  $N_{\mathbb{R}} = \mathbb{R}$ , so that  $M_{\mathbb{R}} = \mathbb{R}$ . Set  $C = [0, 1]$  and  $g_i = 0$ ,  $i = 1, 2$ . Then  $g_i^{\vee}(u) = \min(0, u)$  for  $u \in \mathbb{R}$ . Consider the sequence of measures

$$\mu_l = \frac{l-1}{l} \delta_0 + \frac{1}{l} \delta_{-l}, \quad l \geq 1,$$

where  $\delta_0$  and  $\delta_{-l}$  are the Dirac measures at the points 0 and  $-l$ , respectively. This sequence converges to  $\delta_0$ . However,  $\Phi(\mu_l) = -1$  for all  $l$  and  $\Phi(\delta_0) = 0$ .

Nevertheless, we have the following result.

**Proposition 3.7.** *The function  $\Phi$  is upper semicontinuous.*

To prove this proposition, we need the following lemma.

**Lemma 3.8.** *Let  $\phi: N_{\mathbb{R}} \rightarrow \mathbb{R}$  be a continuous function. If  $\phi$  is bounded above (respectively below), then the map  $\mathcal{P} \rightarrow \mathbb{R} \cup \{-\infty\}$  (respectively  $\mathcal{P} \rightarrow \mathbb{R} \cup \{\infty\}$ ) given by*

$$\mu \mapsto \int \phi d\mu$$

*is upper semicontinuous (respectively lower semicontinuous).*

*Proof.* We prove the case of a function bounded above, the other case being analogous. Let  $\mu \in \mathcal{P}$  and  $\varepsilon > 0$  be given and, for  $l \geq 1$ , put

$$\phi_l(u) = \max(\phi(u), -l).$$

The sequence of functions  $(\phi_l)_{l \geq 1}$  is monotone and converges pointwise to  $\phi$ . So Lebesgue's monotone convergence theorem implies that there is  $l_0 \geq 1$  such that

$$\int \phi_{l_0} \, d\mu \leq \int \phi \, d\mu + \varepsilon.$$

Let  $(\mu_l)_{l \geq 1}$  be a sequence in  $\mathcal{P}$  converging to  $\mu$ . Since  $\phi_{l_0} \in \mathcal{C}_b(N_{\mathbb{R}})$ , there exists  $l_1 \geq 1$  such that, for  $l \geq l_1$ ,

$$\int \phi \, d\mu_l \leq \int \phi_{l_0} \, d\mu_l \leq \int \phi_{l_0} \, d\mu + \varepsilon \leq \int \phi \, d\mu + 2\varepsilon.$$

Since  $\varepsilon$  is arbitrary,  $\limsup_{l \rightarrow \infty} \int \phi \, d\mu_l \leq \int \phi \, d\mu$ , proving the lemma.  $\square$

*Proof of Proposition 3.7.* Set  $\phi_i = g_i^\vee$ ,  $i = 1, 2$  for short. Fix  $\mu_0 \in \mathcal{E}$  and set  $u_0 = -\mathbb{E}[\mu_0] \in N_{\mathbb{R}}$ . Take  $x \in \partial\phi_2(u_0) \subset M_{\mathbb{R}}$ . Then, for all  $u \in N_{\mathbb{R}}$ ,

$$\langle x, u - u_0 \rangle \geq \phi_2(u) - \phi_2(u_0).$$

Let  $\mu \in \mathcal{E}$ . It follows from this inequality that

$$\begin{aligned} \Phi(\mu) - \Phi(\mu_0) &= \int \phi_1 \, d\mu + \phi_2(-\mathbb{E}[\mu]) - \int \phi_1 \, d\mu_0 - \phi_2(-\mathbb{E}[\mu_0]) \\ &\leq \int \phi_1 \, d(\mu - \mu_0) - \langle \mathbb{E}[\mu] - \mathbb{E}[\mu_0], x \rangle \\ &\leq \int \phi_1 \, d(\mu - \mu_0) - \int \langle u, x \rangle \, d(\mu - \mu_0) \\ &\leq \int \phi \, d(\mu - \mu_0) \end{aligned}$$

with  $\phi = \phi_1 - x$ . Hence

$$\Phi(\mu) \leq \Phi(\mu_0) - \int \phi \, d\mu_0 + \int \phi \, d\mu. \quad (3.7)$$

Since  $x$  belongs to  $\partial\phi_2(u_0)$  and  $\partial\phi_2(u_0) \subset \text{stab}(\phi_2) = \text{stab}(\phi_1) = C$ , the function  $\phi$  is bounded above. By Lemma 3.8, the right-hand side of (3.7) is upper semicontinuous. The inequality (3.7) is an equality for  $\mu = \mu_0$ . Hence  $\Phi$  is upper semicontinuous at  $\mu_0$ , as stated.  $\square$

**Proposition 3.9.** *The function  $\Phi$  is nonpositive, and vanishes for  $\mu \in \mathcal{E}$  if and only if*

$$\text{supp}(\mu) \subset F(g_1, g_2) \quad \text{and} \quad \mathbb{E}[\mu] \in B(g_1, g_2), \quad (3.8)$$

with  $B(g_1, g_2)$  and  $F(g_1, g_2)$  as in Definition 3.2.

*Proof.* Let notation be as in Lemma 3.1 and fix a point  $x \in \text{ri}(C_{\max})$ . For short put

$$A_i = \partial g_i(x), \quad i = 1, 2, \quad B = B(g_1, g_2), \quad F = F(g_1, g_2).$$

By Lemma 3.1 (1), the sets  $A_1$  and  $A_2$  do not depend on the choice of the point  $x \in \text{ri}(C_{\max})$ . Let  $\hat{\phi}_i$  be as in (3.2). For every  $\mu \in \mathcal{E}$  we can write  $\Phi(\mu)$  in terms of the functions  $\hat{\phi}_i$  as

$$\Phi(\mu) = \int \hat{\phi}_1 \, d\mu + \hat{\phi}_2(-\mathbb{E}[\mu]). \quad (3.9)$$

By Lemma 3.1(4), the functions  $\hat{\phi}_i$  are nonpositive and vanish precisely on the sets  $A_i$ . It follows from (3.9) that  $\Phi$  is nonpositive and vanishes for every  $\mu \in \mathcal{E}$  satisfying (3.8).

Conversely, let  $\mu \in \mathcal{E}$  such that  $\Phi(\mu) = 0$ . Since both  $\hat{\phi}_1$  and  $\hat{\phi}_2$  are nonpositive, the equality (3.9) also implies that

$$\int \hat{\phi}_1 \, d\mu = 0 \quad \text{and} \quad \hat{\phi}_2(-\mathbb{E}[\mu]) = 0.$$

Therefore  $\text{supp}(\mu) \subset A_1$  and  $-\mathbf{E}[\mu] \in A_2$ . By Proposition 3.5(1),  $\mathbf{E}[\mu]$  belongs to  $\text{conv}(\text{supp}(\mu))$ . Since  $A_1$  is a convex set that contains  $\text{supp}(\mu)$ , we deduce  $\mathbf{E}[\mu] \in A_1$  and so

$$\mathbf{E}[\mu] \in A_1 \cap (-A_2) = B,$$

which gives the second condition in (3.8).

We next prove that the first condition in (3.8) is satisfied. Write  $\theta = \mu(F)$ , so that  $0 \leq \theta \leq 1$  and  $\mu(A_1 \setminus F) = 1 - \theta$ . We want to show  $\theta = 1$ , thus we assume the contrary, namely  $\theta < 1$ . This implies that  $F$  is a proper face of  $A_1$ . We put

$$u_2 = \frac{1}{1 - \theta} \int_{A_1 \setminus F} u \, d\mu \in A_1 \setminus F.$$

If  $\theta = 0$ , then  $\mathbf{E}[\mu] = u_2$  and so  $\mathbf{E}[\mu] \in A_1 \setminus F$ , contradicting the fact that  $\mathbf{E}[\mu] \in B \subset F$ . Suppose that  $0 < \theta < 1$  and set

$$u_1 = \frac{1}{\theta} \int_F u \, d\mu \in F.$$

Therefore

$$\mathbf{E}[\mu] = \theta u_1 + (1 - \theta)u_2 \in \text{ri}(\overline{u_1 u_2}),$$

the relative interior of the segment  $\overline{u_1 u_2}$ . Since  $\mathbf{E}[\mu]$  is in  $B$  and hence in  $F$ , we have  $\text{ri}(\overline{u_1 u_2}) \cap F \neq \emptyset$ . Moreover, the whole segment is contained in  $A_1$ , and  $F$  is a face of  $A_1$ . We deduce that this segment is contained in  $F$ . Therefore  $u_2 \in F$ , contradicting the fact that  $u_2 \in A_1 \setminus F$ . We conclude that  $\theta = 1$  and so  $\text{supp}(\mu) \subset F$  since  $F$  is closed. This proves the first condition and completes the proof.  $\square$

The function  $\Phi$  satisfies also the following property.

**Lemma 3.10.** *There are constants  $c_1 \geq 0$  and  $c_2 > 0$  such that, for all  $\mu \in \mathcal{E}$ ,*

$$\Phi(\mu) \leq c_1 - c_2 \int \|u\| \, d\mu.$$

*Proof.* Let  $\Psi$  be the support function of  $C$ , which is the function on  $N_{\mathbb{R}}$  given by

$$\Psi(u) = \min_{y \in C} \langle u, y \rangle.$$

Put  $c_1 = 4 \max_{y \in C} (|g_1(y)|, |g_2(y)|)$ . It follows from their definition that the functions  $\phi_i = g_i^{\vee}$  verify, for  $u \in N_{\mathbb{R}}$ ,

$$\max(\phi_1(u), \phi_2(u)) \leq \Psi(u) + \frac{c_1}{4}. \quad (3.10)$$

Let  $x$  be a point in the interior of  $C$ . On  $M_{\mathbb{R}}$ , we consider the norm induced by the chosen norm  $\|\cdot\|$  in  $N_{\mathbb{R}}$ . Since  $x$  is in the interior of  $C$ , we can find a constant  $c_2 > 0$  such that  $\mathcal{B}(x, c_2)$ , the closed ball of center  $x$  and radius  $c_2$ , is contained in  $C$ . Then

$$\Psi(u) \leq \min_{y \in \mathcal{B}(x, c_2)} \langle u, y \rangle = \langle u, x \rangle - c_2 \|u\|. \quad (3.11)$$

Since  $x \in C = \text{stab}(\Psi)$ , we have  $(\Psi - x)(u) \leq 0$ . By (3.10) and (3.11),

$$\begin{aligned} \Phi(\mu) &= \int \phi_1(u) \, d\mu + \phi_2(-\mathbf{E}[\mu]) + \max_{y \in C} (g_1(y) + g_2(y)) \\ &\leq c_1 + \int \Psi(u) \, d\mu + \Psi(-\mathbf{E}[\mu]) \\ &\leq c_1 + \int (\Psi - x)(u) \, d\mu + (\Psi - x)(-\mathbf{E}[\mu]) \\ &\leq c_1 - c_2 \int \|u\| \, d\mu, \end{aligned}$$

as stated.  $\square$

**Proposition 3.11.** *Let  $(\mu_l)_{l \in I}$  be a net of measures in  $\mathcal{E}$  such that*

$$\lim_l \Phi(\mu_l) = 0.$$

*Then  $(\mu_l)_{l \in I}$  has at least one cluster point in  $\mathcal{P}$ , and every such cluster point  $\mu$  lies in  $\mathcal{E}$  and satisfies*

$$\text{supp}(\mu) \subset F(g_1, g_2) \quad \text{and} \quad E[\mu] \in B(g_1, g_2).$$

*Proof.* Replacing  $(\mu_l)_{l \in I}$  by a subnet if necessary, we assume that  $\Phi(\mu_l) \geq -1$  for all  $l \in I$ . Let  $c_1, c_2$  be the constants of Lemma 3.10 and set  $K = (c_1 + 1)/c_2 > 0$ . This lemma implies that each  $\mu_l$  is in the subset of  $\mathcal{E}$  given by

$$\left\{ \mu \in \mathcal{E} \mid \int \|u\| \, d\mu(u) \leq K \right\}.$$

This subset is compact thanks to Prokhorov's theorem [Bou69, *Théorème 5.5.1*], and it is metrizable because  $\mathcal{P}$  is. Hence, the net  $(\mu_l)_{l \in I}$  has at least one cluster point, and every such cluster point  $\mu$  lies in  $\mathcal{E}$ , proving the first statement.

To prove the last statement, let  $(\mu_k)_{k \in I'}$  be a subnet converging to  $\mu$ . By Proposition 3.7, the function  $\Phi$  is upper-semicontinuous and so

$$\Phi(\mu) \geq \limsup_k \Phi(\mu_k) = 0.$$

Hence  $\Phi(\mu) = 0$ , and the statement follows from Proposition 3.9.  $\square$

As we have seen in Example 3.6, the function  $\Phi$  is not continuous. We now consider another topology on  $\mathcal{E}$  with respect to which the function  $\Phi$  is continuous.

Given  $\mu, \mu' \in \mathcal{P}$ , denote by  $\Gamma(\mu, \mu')$  the set of probability measures on  $N_{\mathbb{R}} \times N_{\mathbb{R}}$  with marginals  $\mu$  and  $\mu'$ . That is, a probability measure  $\nu$  on  $N_{\mathbb{R}} \times N_{\mathbb{R}}$  belongs to  $\Gamma(\mu, \mu')$  if and only if

$$(p_1)_* \nu = \mu, \quad (p_2)_* \nu = \mu',$$

where  $p_i$  is the projection of  $N_{\mathbb{R}} \times N_{\mathbb{R}}$  onto its  $i$ -th factor, and  $(p_i)_*$  the direct image of measures.

**Definition 3.12.** The *Kantorovich–Rubinstein distance* (or *Wasserstein distance of order 1*) on  $\mathcal{E}$  is defined, for  $\mu, \mu' \in \mathcal{E}$ , by

$$W(\mu, \mu') = \inf_{\nu \in \Gamma(\mu, \mu')} \int \|u - u'\| \, d\nu(u, u').$$

The quantity  $W(\mu, \mu')$  satisfies the axioms of a distance and is finite when  $\mu, \mu' \in \mathcal{E}$  [Vil09, pages 94–5]. The *Kantorovich–Rubinstein topology* (or *KR-topology* for short) of  $\mathcal{E}$  is the topology induced by this distance.

The finiteness of  $W(\mu, \mu')$  for  $\mu$  and  $\mu'$  in  $\mathcal{E}$ , can be argued as follows. The product measure  $\mu \times \mu'$  is in  $\Gamma(\mu, \mu')$ , and we have

$$W(\mu, \mu') \leq \int \|u - u'\| \, d(\mu \times \mu')(u, u') \leq \int \|u\| \, d\mu(u) + \int \|u'\| \, d\mu'(u') < \infty.$$

For a Lipschitz continuous function  $\psi: N_{\mathbb{R}} \rightarrow \mathbb{R}$ , denote by  $\text{Lip}(\psi)$  its *Lipschitz constant*, given by

$$\text{Lip}(\psi) = \sup_{u \neq u'} \frac{|\psi(u) - \psi(u')|}{\|u - u'\|}.$$

Lipschitz constants and the Kantorovich–Rubinstein distance are related by the *duality formula*: for  $\mu, \mu' \in \mathcal{E}$  and a Lipschitz continuous function  $\psi: N_{\mathbb{R}} \rightarrow \mathbb{R}$ , we have

$$\left| \int \psi \, d\mu - \int \psi \, d\mu' \right| \leq \text{Lip}(\psi) W(\mu, \mu'), \quad (3.12)$$

see for instance [Vil09, Remark 6.5].

**Remark 3.13.** By [Vil09, Theorem 6.9], the KR-topology agrees with the weak-\* topology on  $\mathcal{E}$  with respect to the space of continuous functions  $\varphi: N_{\mathbb{R}} \rightarrow \mathbb{R}$  such that

$$|\varphi(u)| \leq c(1 + \|u\|)$$

for a  $c \in \mathbb{R}$  and all  $u \in N_{\mathbb{R}}$ . In particular, the KR-topology is stronger than the topology of  $\mathcal{E}$  induced by that of  $\mathcal{P}$  as in Definition 3.4.

**Proposition 3.14.** *The function  $\Phi$  is continuous with respect to the KR-topology. In particular, if  $(\mu_l)_{l \in I}$  is a net of measures in  $\mathcal{E}$  that converges to a measure  $\mu \in \mathcal{E}$  with respect to the KR-topology and*

$$\text{supp}(\mu) \subset F(g_1, g_2) \quad \text{and} \quad E[\mu] \in B(g_1, g_2),$$

then  $\lim_l \Phi(\mu_l) = 0$ .

*Proof.* Let  $(\mu_l)_{l \in I}$  be a net on  $\mathcal{E}$  that converges to a measure  $\mu \in \mathcal{E}$  with respect to the KR-topology. By Remark 3.13,

$$\lim_l \int g_1^\vee d\mu_l = \int g_1^\vee d\mu \quad \text{and} \quad \lim_l g_2^\vee(-E[\mu_l]) = g_2^\vee(-E[\mu]).$$

Therefore  $\lim_l \Phi(\mu_l) = \Phi(\mu)$  and so  $\Phi$  is continuous, proving the first statement. The second statement follows from the first one and Proposition 3.9.  $\square$

We also need the following easy result. We include it here for the lack of a suitable reference.

**Lemma 3.15.** *Let  $E_i \subset N_{\mathbb{R}}$ ,  $i = 1, \dots, r$ , be convex subsets and  $E = E_1 + \dots + E_r$  their Minkowski sum. For a point  $u_0 \in E$ , the following conditions are equivalent:*

- (1) *the point  $u_0$  is a vertex of  $E$ ;*
- (2) *the equation  $u_0 = \sum_i z_i$  with  $z_i \in E_i$  has a unique solution and, for  $i = 1, \dots, r$ , the point  $z_i$  in this solution is a vertex of  $E_i$ .*

*Proof.* First assume that  $u_0$  is a vertex of  $E$ . Suppose that the equation  $u_0 = \sum_i z_i$ ,  $z_i \in E_i$ , has two different solutions, namely  $u_0 = \sum_i z'_i$  and  $u_0 = \sum_i z''_i$  with  $z'_{i_0} \neq z''_{i_0}$  for some  $i_0 \in \{1, \dots, r\}$ . Then both points

$$u_1 = \sum_{i \neq i_0} z'_i + z''_{i_0} \quad \text{and} \quad u_2 = \sum_{i \neq i_0} z''_i + z'_{i_0}$$

belong to  $E$ , they are different and satisfy  $u_0 = \frac{1}{2}(u_1 + u_2)$ , contradicting the fact that  $u_0$  is a vertex of  $E$ . Hence the equation  $u_0 = \sum_i z_i$  has a unique solution with  $z_i \in E_i$ .

Now suppose that  $z_{i_0}$  is not a vertex of  $E_{i_0}$  for some  $i_0 \in \{1, \dots, r\}$ . Then we can write  $z_{i_0} = \frac{1}{2}(z'_{i_0} + z''_{i_0})$  with  $z'_{i_0} \neq z''_{i_0}$  both in  $E_{i_0}$ . Hence the points

$$u_1 = \sum_{i \neq i_0} z_i + z'_{i_0} \quad \text{and} \quad u_2 = \sum_{i \neq i_0} z_i + z''_{i_0}$$

are different, belong to  $E$  and  $u_0 = \frac{1}{2}(u_1 + u_2)$ , contradicting the assumption that  $u_0$  is a vertex of  $E$ . Thus we have proved that (1) implies (2).

Assume now that the statement (2) is true but  $u_0$  is not a vertex of  $E$ . Then there are two different points  $u_1, u_2 \in E$  with  $u_0 = \frac{1}{2}(u_1 + u_2)$ . Since  $E$  is the Minkowski sum of the sets  $E_i$ , we can write

$$u_0 = \sum_i z_i, \quad u_1 = \sum_i z'_i \quad \text{and} \quad u_2 = \sum_i z''_i.$$

The equation  $u_0 = \sum_i z_i$  has a unique solution and so  $z_i = \frac{1}{2}(z'_i + z''_i)$  for all  $i$ . Since  $z_i$  is a vertex of  $E_i$ , this implies  $z'_i = z''_i$ . Therefore  $u_1 = u_2$ , contradicting the assumptions and thus proving that (2) implies (1).  $\square$

## 4. MODULUS DISTRIBUTION

In this section, we study the asymptotic modulus distribution of the Galois orbits of nets of points of small height in toric varieties. Our approach is based on the techniques developed in the series of papers [BPS14, BMPS16, BPS15]. These techniques are well-suited for the study of toric metrics and their associated height functions. In the sequel, we recall the basic constructions and results.

Let  $\mathbb{K}$  be a global field and  $\mathbb{T} \simeq \mathbb{G}_{m,\mathbb{K}}^n$  a split torus of dimension  $n$  over  $\mathbb{K}$ . Let

$$N = \text{Hom}(\mathbb{G}_{m,\mathbb{K}}, \mathbb{T}) \quad \text{and} \quad M = \text{Hom}(\mathbb{T}, \mathbb{G}_{m,\mathbb{K}}) = N^\vee$$

be the lattices of cocharacters and of characters of  $\mathbb{T}$ , respectively, and write  $N_{\mathbb{R}} = N \otimes \mathbb{R}$  and  $M_{\mathbb{R}} = M \otimes \mathbb{R}$ . We also fix an auxiliary norm  $\|\cdot\|$  on  $N_{\mathbb{R}}$ .

Let  $v \in \mathfrak{M}_{\mathbb{K}}$ . We denote by  $\mathbb{T}_v^{\text{an}}$  the  $v$ -adic analytification of  $\mathbb{T}$  and by  $\mathbb{S}_v$  its compact subtorus. In the Archimedean case,  $\mathbb{S}_v$  is isomorphic to  $(S^1)^n$  whereas, in the non-Archimedean case, it is a compact analytic group, see [BPS14, § 4.2] for a description. Moreover, there is a map  $\text{val}_v: \mathbb{T}_v^{\text{an}} \rightarrow N_{\mathbb{R}}$ , defined, in a given splitting, by

$$\text{val}_v(x_1, \dots, x_n) = (-\log |x_1|_v, \dots, -\log |x_n|_v). \quad (4.1)$$

This map does not depend on the choice of the splitting, and the compact torus  $\mathbb{S}_v$  coincides with its fiber over the point  $0 \in N_{\mathbb{R}}$ .

Let  $X$  be a proper toric variety over  $\mathbb{K}$  with torus  $\mathbb{T}$ , described by a complete fan  $\Sigma$  on  $N_{\mathbb{R}}$ . To each cone  $\sigma \in \Sigma$  corresponds an affine toric variety  $X_\sigma$ , which is an open subset of  $X$ , and an orbit  $O(\sigma)$  of the action of  $\mathbb{T}$  on  $X$ . The affine toric variety corresponding to the cone  $\sigma = \{0\}$  is the *principal open subset*  $X_0$ . It coincides with the orbit  $O(0)$  and is canonically isomorphic to the torus  $\mathbb{T}$ .

An  $\mathbb{R}$ -divisor  $D$  on  $X$  is *toric* if it is invariant under the action of  $\mathbb{T}$ . Such an  $\mathbb{R}$ -divisor defines a *virtual support function* on  $\Sigma$ , that is, a function

$$\Psi_D: N_{\mathbb{R}} \longrightarrow \mathbb{R}$$

whose restriction to each cone of the fan  $\Sigma$  is linear. We also associate to  $D$  the subset of  $M_{\mathbb{R}}$  given by

$$\Delta_D = \text{stab}(\Psi_D) = \{x \in M_{\mathbb{R}} \mid x \geq \Psi_D\}.$$

If  $D$  is pseudo-effective, then  $\Delta_D$  is a polytope and, otherwise, it is the empty set. Properties of the  $\mathbb{R}$ -divisor  $D$  can be read off from its associated virtual support function and polytope. In particular,  $D$  is nef if and only if  $\Psi_D$  is concave, and  $D$  is big if and only if  $\Delta_D$  has nonempty interior.

A quasi-algebraic metrized divisor  $\overline{D} = (D, (\|\cdot\|_v)_{v \in \mathfrak{M}_{\mathbb{K}}})$  on  $X$  is *toric* if and only if the  $v$ -adic metric  $\|\cdot\|_v$  is invariant with respect to the action of  $\mathbb{S}_v$ , for all  $v$ . Such a toric metrized  $\mathbb{R}$ -divisor on  $X$  defines a family of continuous functions  $\psi_{\overline{D},v}: N_{\mathbb{R}} \rightarrow \mathbb{R}$  indexed by the places of  $\mathbb{K}$ . For each  $v \in \mathfrak{M}_{\mathbb{K}}$ , this function is given, for  $p \in \mathbb{T}_v^{\text{an}}$ , by

$$\psi_{\overline{D},v}(\text{val}_v(p)) = \log \|s_D(p)\|_v, \quad (4.2)$$

where  $s_D$  is the canonical rational  $\mathbb{R}$ -section of  $D$  as in [BMPS16, §3]. This adelic family of functions satisfies that  $|\psi_{\overline{D},v} - \Psi_D|$  is bounded for all  $v$ , and that  $\psi_{\overline{D},v} = \Psi_D$  for all  $v$  except for a finite number. In particular, the stability set of each  $\psi_{\overline{D},v}$  coincides with  $\Delta_D$ .

For each  $v \in \mathfrak{M}_{\mathbb{K}}$ , we also consider the  *$v$ -adic roof function*  $\vartheta_{\overline{D},v}: \Delta_D \rightarrow \mathbb{R}$ , which is given by

$$\vartheta_{\overline{D},v}(x) = \psi_{\overline{D},v}^\vee(x) = \inf_{u \in N_{\mathbb{R}}} (\langle u, x \rangle - \psi_{\overline{D},v}(u)).$$

This is an adelic family of continuous concave functions on  $\Delta_D$  that are zero except for a finite number of places. The *global roof function*  $\vartheta_{\overline{D}}: \Delta_D \rightarrow \mathbb{R}$  is the weighted sum

$$\vartheta_{\overline{D}} = \sum_{v \in \mathfrak{M}_{\mathbb{K}}} n_v \vartheta_{\overline{D},v}.$$

The essential minimum of  $X$  with respect to  $\overline{D}$  defined in (2.6) can be computed as the maximum of its roof function [BPS15, Theorem A], that is

$$\mu_{\overline{D}}^{\text{ess}}(X) = \max_{x \in \Delta_D} \vartheta_{\overline{D}}(x). \quad (4.3)$$

**Example 4.1.** Let  $X$  be a proper toric variety over  $\mathbb{K}$  and  $D$  a toric  $\mathbb{R}$ -divisor on  $X$ . The *canonical metric* on  $D$  is the metric characterized by the fact that, for each  $v \in \mathfrak{M}_{\mathbb{K}}$  and  $p \in \mathbb{T}_v^{\text{an}}$ ,

$$\log \|s_D(p)\|_{\text{can},v} = \Psi_D(\text{val}_v(p)),$$

see [BPS14, Proposition-Definition 4.3.15]. We denote this toric metrized  $\mathbb{R}$ -divisor by  $\overline{D}^{\text{can}}$ . For all  $v \in \mathfrak{M}_{\mathbb{K}}$ ,

$$\psi_{\overline{D}^{\text{can}},v} = \Psi_D \quad \text{and} \quad \vartheta_{\overline{D}^{\text{can}},v} = 0.$$

In particular,  $\vartheta_{\overline{D}^{\text{can}}} = 0$  and  $\mu_{\overline{D}^{\text{can}}}^{\text{ess}}(X) = 0$ .

Given a semipositive toric metrized  $\mathbb{R}$ -divisor  $\overline{D}$  over  $D$ , its associated metric functions are concave. Conversely, every adelic family of concave continuous functions  $\psi_v: N_{\mathbb{R}} \rightarrow \mathbb{R}$ ,  $v \in \mathfrak{M}_{\mathbb{K}}$ , with  $|\psi_v - \Psi_D|$  bounded for all  $v$  and such that  $\psi_{\overline{D},v} = \Psi_D$  for all  $v$  except for a finite number, corresponds to a semipositive toric metrized  $\mathbb{R}$ -divisor over  $D$  [BMPS16, Proposition 4.19(1)]. For instance, a canonical toric metrized  $\mathbb{R}$ -divisor  $\overline{D}^{\text{can}}$  is semipositive if and only if  $\Psi_D$  is concave, which is equivalent to the condition that  $D$  is nef.

For the rest of this section, we suppose that  $X$  is a proper toric variety over the global field  $\mathbb{K}$  with torus  $\mathbb{T}$ , and that  $\overline{D}$  is a semipositive toric metrized  $\mathbb{R}$ -divisor with  $D$  big.

We also fix the notation below. Recall from §3 that  $\mathcal{P}$  denotes the space of probability measures on  $N_{\mathbb{R}}$  endowed with the weak- $*$  topology with respect to the space  $\mathcal{C}_b(N_{\mathbb{R}})$ , and that  $\mathcal{E}$  denotes the subspace of probability measures with finite first moment.

**Notation 4.2.** Let  $v \in \mathfrak{M}_{\mathbb{K}}$ . We denote by  $g_{i,v}$ ,  $i = 1, 2$ , the concave functions on  $\Delta_D$  given by

$$g_{1,v} = \vartheta_{\overline{D},v} \quad \text{and} \quad g_{2,v} = \sum_{w \in \mathfrak{M}_{\mathbb{K}} \setminus \{v\}} \frac{n_w}{n_v} \vartheta_{\overline{D},w}.$$

Thus  $\vartheta_{\overline{D}} = n_v(g_{1,v} + g_{2,v})$ . We consider the convex subsets of  $N_{\mathbb{R}}$  given by Definition 3.2

$$B_v = B(g_{1,v}, g_{2,v}), \quad F_v = F(g_{1,v}, g_{2,v}) \quad (4.4)$$

and we write

$$A_v = \partial g_{1,v}(x)$$

for any  $x$  in the relative interior of the set  $\Delta_{D,\max}$  where  $\vartheta_{\overline{D}}$  attains its maximum. By Lemma 3.1 (1)  $A_v$  does not depend on the choice of  $x \in \text{ri}(\Delta_{D,\max})$ . Thus  $F_v$  is the minimal face of  $A_v$  containing  $B_v$ . We also denote  $\Phi_v$  the function on  $\mathcal{E}$  given by Definition (3.6) applied to the set  $C = \Delta_D$  and the functions  $g_{i,v}$ ,  $i = 1, 2$ .

Given  $v \in \mathfrak{M}_{\mathbb{K}}$  and a point  $p \in X_0(\overline{\mathbb{K}})$ , we consider the discrete probability measure on  $N_{\mathbb{R}}$  defined by

$$\nu_{p,v} = (\text{val}_v)_* \mu_{p,v},$$

where  $\mu_{p,v}$  is the uniform discrete probability measure on  $X_v^{\text{an}}$  supported on the  $v$ -adic Galois orbit of  $p$  as in (2.4). This probability measure on  $N_{\mathbb{R}}$  gives the modulus distribution of the  $v$ -adic Galois orbit of the point  $p$ . The next result characterizes the limit behavior of this modulus distribution for nets of points of small height.

**Theorem 4.3.** *Let notation and hypotheses be as above. For each  $v \in \mathfrak{M}_{\mathbb{K}}$  and every  $\overline{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points in the principal open subset  $X_0$ , the net  $(\nu_{p_l, v})_{l \in I}$  of measures in  $\mathcal{P}$  has at least one cluster point. Every such cluster point  $\nu_v$  lies in  $\mathcal{E}$  and satisfies*

$$\text{supp}(\nu_v) \subset F_v \quad \text{and} \quad \mathbb{E}[\nu_v] \in B_v. \quad (4.5)$$

The proof of Theorem 4.3 is given below, after a definition and an auxiliary result.

**Definition 4.4.** A centered adelic measure  $\nu$  on  $N_{\mathbb{R}}$  is a collection of measures  $\nu_v \in \mathcal{E}$ ,  $v \in \mathfrak{M}_{\mathbb{K}}$ , such that  $\nu_v = \delta_0$ , the Dirac measure at the point  $0 \in N_{\mathbb{R}}$ , for all but a finite number of places  $v$ , and such that

$$\sum_{v \in \mathfrak{M}_{\mathbb{K}}} n_v \mathbb{E}[\nu_v] = 0. \quad (4.6)$$

We denote by  $\mathcal{H}_{\mathbb{K}}$  the set of all centered adelic measures on  $N_{\mathbb{R}}$ .

We introduce the function  $\eta_{\overline{D}}: \mathcal{H}_{\mathbb{K}} \rightarrow \mathbb{R}$  defined by

$$\eta_{\overline{D}}(\nu) = - \sum_{v \in \mathfrak{M}_{\mathbb{K}}} n_v \int \psi_{\overline{D}, v} d\nu_v. \quad (4.7)$$

This function extends the notion of height of points to the space  $\mathcal{H}_{\mathbb{K}}$ . Indeed, for  $p \in X_0(\overline{\mathbb{K}})$ , the collection

$$\nu_p = (\nu_{p, v})_{v \in \mathfrak{M}_{\mathbb{K}}} \quad (4.8)$$

is a centered adelic measure on  $N_{\mathbb{R}}$ , because of the product formula in Proposition 2.1(2). Moreover, the canonical  $\mathbb{R}$ -section  $s_D$  does not vanish at  $p$  and, by Proposition 2.3 and (4.2),

$$\begin{aligned} h_{\overline{D}}(p) &= - \sum_v \frac{n_v}{\#\text{Gal}(p)_v} \sum_{q \in \text{Gal}(p)_v} \psi_{\overline{D}, v}(\text{val}_v(q)) \\ &= - \sum_v n_v \int \psi_{\overline{D}, v} d\nu_{p, v} = \eta_{\overline{D}}(\nu_p). \end{aligned} \quad (4.9)$$

**Lemma 4.5.** *For every centered adelic measure  $\nu = (\nu_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ ,*

$$\max_{v \in \mathfrak{M}_{\mathbb{K}}} -n_v \Phi_v(\nu_v) \leq \eta_{\overline{D}}(\nu) - \mu_{\overline{D}}^{\text{ess}}(X) \leq \sum_{v \in \mathfrak{M}_{\mathbb{K}}} -n_v \Phi_v(\nu_v). \quad (4.10)$$

*In particular, for  $p \in X_0(\overline{\mathbb{K}})$ ,*

$$\max_{v \in \mathfrak{M}_{\mathbb{K}}} -n_v \Phi_v(\nu_{p, v}) \leq h_{\overline{D}}(p) - \mu_{\overline{D}}^{\text{ess}}(X) \leq \sum_{v \in \mathfrak{M}_{\mathbb{K}}} -n_v \Phi_v(\nu_{p, v}). \quad (4.11)$$

*Proof.* Let  $\Delta_{D, \max}$  be the set of points maximizing the roof function  $\vartheta_{\overline{D}}$  and choose  $x \in \Delta_{D, \max}$ . For each  $v \in \mathfrak{M}_{\mathbb{K}}$ , let  $\widehat{\phi}_{i, v}: N_{\mathbb{R}} \rightarrow \mathbb{R}$ ,  $i = 1, 2$ , be the function defined by

$$\widehat{\phi}_{i, v}(u) = g_{i, v}^{\vee}(u) - \langle x, u \rangle + g_{i, v}(x),$$

where  $g_{i, v}$  denotes the concave function on  $\Delta_D$  in Notation 4.2 and  $g_{i, v}^{\vee}$  its Legendre dual as in (3.1).



Note that  $\psi_{\overline{D},v} = g_{1,v}^\vee$ . Using (4.6) and (4.3), we deduce that

$$-\sum_v n_v \int \psi_{\overline{D},v} d\nu_v = \vartheta_{\overline{D}}(x) - \sum_v n_v \int \widehat{\phi}_{1,v} d\nu_v = \mu_{\overline{D}}^{\text{ess}}(X) - \sum_v n_v \int \widehat{\phi}_{1,v} d\nu_v.$$

Thus

$$\eta_{\overline{D}}(\nu) - \mu_{\overline{D}}^{\text{ess}}(X) = -\sum_v n_v \int \widehat{\phi}_{1,v} d\nu_v. \quad (4.12)$$

For each  $v \in \mathfrak{M}_{\mathbb{K}}$ , we get from the definition of  $\Phi_v$  that

$$\Phi_v(\nu_v) = \int \widehat{\phi}_{1,v} d\nu_v + \widehat{\phi}_{2,v}(-\mathbf{E}[\nu_v]).$$

By Lemma 3.1(4), the functions  $\widehat{\phi}_{i,v}$  are nonpositive and so

$$\Phi_v(\nu_v) \leq \int \widehat{\phi}_{1,v} d\nu_v. \quad (4.13)$$

The second inequality in (4.10) then follows from (4.12) and (4.13).

To prove the first inequality in (4.10), fix  $v \in \mathfrak{M}_{\mathbb{K}}$ . By [BPS14, Propositions 2.3.1(1) and 2.3.3(3)],

$$\widehat{\phi}_{2,v} = \boxplus_{w \neq v} \left( \widehat{\phi}_{1,w} \frac{n_w}{n_v} \right), \quad (4.14)$$

where  $w$  runs over the places of  $\mathbb{K}$  different from  $v$ , the symbol  $\boxplus$  denotes the sup-convolution and, for a concave function  $\psi$  and a nonzero constant  $\lambda$ , the expression  $\psi\lambda$  denotes the right multiplication as in [BPS14, §2.3].

By the equality (4.14), the definitions of the sup-convolution and the right multiplication, and condition (4.6), we deduce

$$\widehat{\phi}_{2,v}(-\mathbf{E}[\nu_v]) \geq \sum_{w \neq v} \frac{n_w}{n_v} \widehat{\phi}_{1,w}(\mathbf{E}[\nu_w]). \quad (4.15)$$

By the concavity of  $\widehat{\phi}_{1,w}$  and Jensen's inequality,  $\int \widehat{\phi}_{1,w} d\nu_w \leq \widehat{\phi}_{1,w}(\mathbf{E}[\nu_w])$  for all  $w \in \mathfrak{M}_{\mathbb{K}}$ . Therefore, by (4.12) and (4.15),

$$\begin{aligned} \eta_{\overline{D}}(\nu) - \mu_{\overline{D}}^{\text{ess}}(X) &\geq -n_v \left( \int \widehat{\phi}_{1,v} d\nu_v + \sum_{w \neq v} \frac{n_w}{n_v} \widehat{\phi}_{1,w}(\mathbf{E}[\nu_w]) \right) \\ &\geq -n_v \left( \int \widehat{\phi}_{1,v} d\nu_v + \widehat{\phi}_{2,v}(-\mathbf{E}[\nu_v]) \right) = -n_v \Phi_v(\nu_v), \end{aligned}$$

which proves the first inequality and completes the proof of (4.10). The inequalities in (4.11) follow directly from (4.10) and (4.9).  $\square$

*Proof of Theorem 4.3.* Let  $v \in \mathfrak{M}_{\mathbb{K}}$  and let  $\Phi_v: \mathcal{E} \rightarrow \mathbb{R}$  be the function defined by (3.6) with  $g_{1,v}$  and  $g_{2,v}$  as in Notation 4.2. Since the net of points  $(p_l)_{l \in I}$  is  $\overline{D}$ -small,

$$\lim_l h_{\overline{D}}(p_l) = \mu_{\overline{D}}^{\text{ess}}(X).$$

By Proposition 3.9,  $\Phi_v$  is nonpositive, and so we deduce from Lemma 4.5 that

$$\lim_l \Phi_v(\nu_{p_l,v}) = 0.$$

The theorem is then a direct consequence of Proposition 3.11.  $\square$

To state a partial converse of Theorem 4.3, we need a further definition.

**Definition 4.6.** The *adelic Kantorovich–Rubinstein distance*  $W_{\mathbb{K}}$  on  $\mathcal{H}_{\mathbb{K}}$  is defined, for  $\nu = (\nu_v)_v, \nu' = (\nu'_v)_v \in \mathcal{H}_{\mathbb{K}}$ , by

$$W_{\mathbb{K}}(\nu, \nu') = \sum_v n_v W(\nu_v, \nu'_v),$$

where  $W$  denotes the Kantorovich–Rubinstein distance in  $N_{\mathbb{R}}$  as in Definition 3.12. By the definition of  $\mathcal{H}_{\mathbb{K}}$ , there are only a finite number of nonzero terms in this sum.

The topology on  $\mathcal{H}_{\mathbb{K}}$  induced by this distance is called the *adelic KR-topology*.

**Theorem 4.7.** *With notation and hypotheses as before, let  $\nu = (\nu_v)_{v \in \mathfrak{M}_{\mathbb{K}}} \in \mathcal{H}_{\mathbb{K}}$  be a centered adelic measure such that*

$$\text{supp}(\nu_v) \subset F_v \quad \text{and} \quad E[\nu_v] \in B_v$$

for all  $v$ . Then there is a generic  $\bar{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$  such that the net of measures  $(\nu_{p_l})_{l \in I}$  converges to  $\nu$  with respect to the adelic Kantorovich–Rubinstein distance.

The proof of Theorem 4.7 is given below, after some preliminary results. The first result gives the main properties of the function  $\eta_{\bar{D}}$ .

**Lemma 4.8.** *The function  $\eta_{\bar{D}}$  is Lipschitz continuous with respect to  $W_{\mathbb{K}}$ . Moreover, for all  $\nu = (\nu_v)_{v \in \mathfrak{M}_{\mathbb{K}}} \in \mathcal{H}_{\mathbb{K}}$ ,*

$$\eta_{\bar{D}}(\nu) \geq \mu_{\bar{D}}^{\text{ess}}(X), \quad (4.16)$$

with equality if and only if  $\text{supp}(\nu_v) \subset F_v$  and  $E[\nu_v] \in B_v$  for all  $v$ .

*Proof.* Let  $S \subset \mathfrak{M}_{\mathbb{K}}$  be a finite subset such that  $\psi_{\bar{D},v} = \Psi_D$  for all  $v \notin S$ . For  $\nu = (\nu_v)_v, \nu' = (\nu'_v)_v \in \mathcal{H}_{\mathbb{K}}$ ,

$$\begin{aligned} |\eta_{\bar{D}}(\nu) - \eta_{\bar{D}}(\nu')| &\leq \sum_v n_v \left| \int \psi_{\bar{D},v} d\nu_v - \int \psi_{\bar{D},v} d\nu'_v \right| \\ &\leq \sum_v \text{Lip}(\psi_{\bar{D},v}) n_v W(\nu_v, \nu'_v) \leq \left( \max_{x \in \Delta_D} \|x\| \right) W_{\mathbb{K}}(\nu, \nu'). \end{aligned}$$

where the second inequality is given by the duality formula (3.12) and the last by the observation that  $\text{Lip}(\psi_{\bar{D},v}) = \max_{x \in \Delta_D} \|x\|$  for all  $v$ . This proves that  $\eta_{\bar{D}}$  is Lipschitz continuous with respect to  $W_{\mathbb{K}}$ .

As already remarked, the functions  $\Phi_v$  are nonpositive. By Lemma 4.5, this implies the inequality (4.16). From the same result, it follows that the equality holds if and only if  $\Phi_v(\nu_v) = 0$  for all  $v$ . By Proposition 3.9, this holds if and only if  $\text{supp}(\nu_v) \subset F_v$  and  $E[\nu_v] \in B_v$ , completing the proof of the lemma.  $\square$

From this lemma, we deduce as a direct consequence the next characterization of algebraic points in toric varieties realizing the essential minimum.

**Corollary 4.9.** *Let  $p$  be an algebraic point of  $X_0$ . Then  $h_{\bar{D}}(p) = \mu_{\bar{D}}^{\text{ess}}(X)$  if and only if  $\text{supp}(\nu_{p,v}) \subset F_v$  and  $E[\nu_{p,v}] \in B_v$  for all  $v \in \mathfrak{M}_{\mathbb{K}}$ .*

Let  $H_{\mathbb{K}} \subset \bigoplus_{v \in \mathfrak{M}_{\mathbb{K}}} N_{\mathbb{R}}$  be the subspace defined by the equation  $\sum_v n_v u_v = 0$ . By sending the point  $(u_v)_v \in H_{\mathbb{K}}$  to the adelic centered measure  $(\delta_{u_v})_v \in \mathcal{H}_{\mathbb{K}}$ , we identify  $H_{\mathbb{K}}$  with a subspace of  $\mathcal{H}_{\mathbb{K}}$ .

**Corollary 4.10.** *The minimum of the function  $\eta_{\bar{D}}$  is equal to  $\mu_{\bar{D}}^{\text{ess}}(X)$  and it is attained at a point of the subspace  $H_{\mathbb{K}} \subset \mathcal{H}_{\mathbb{K}}$ .*

*Proof.* Let  $x$  be a point where  $\vartheta_{\overline{D}}$  attains its maximum. Since  $0 \in \partial\vartheta_{\overline{D}}(x)$  and  $\partial\vartheta_{\overline{D}}(x) = \sum_v n_v \partial\vartheta_{\overline{D},v}(x)$ , we can find  $\mathbf{u} = (u_v)_v \in H_{\mathbb{K}}$  such that for every  $v$ ,

$$u_v \in \partial\vartheta_{\overline{D},v}(x) \cap \left( -\partial \left( \sum_{w \in \mathfrak{M}_{\mathbb{K}} \setminus \{v\}} \frac{n_w}{n_v} \vartheta_{\overline{D},w} \right) (x) \right) = B_v.$$

The adelic centred measure  $\boldsymbol{\delta} = (\delta_{u_v})_v \in \mathcal{H}_{\mathbb{K}}$  corresponding to  $\mathbf{u} \in H_{\mathbb{K}}$  satisfies  $\text{supp}(\delta_{u_v}) = \{u_v\} \subset F_v$  and  $\mathbb{E}[\delta_{u_v}] = u_v \in B_v$ . Thus, by Lemma 4.8,

$$\mu_{\overline{D}}^{\text{ess}}(X) = \eta_{\overline{D}}(\mathbf{u}) = \min_{\boldsymbol{\nu} \in \mathcal{H}_{\mathbb{K}}} \eta_{\overline{D}}(\boldsymbol{\nu}),$$

as stated.  $\square$

We next show that the measures coming from algebraic points are dense in  $\mathcal{H}_{\mathbb{K}}$ .

**Proposition 4.11.** *For every  $\boldsymbol{\nu} \in \mathcal{H}_{\mathbb{K}}$  there is a generic net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$  such that the net of associated measures  $(\nu_{p_l})_{l \in I}$  as in (4.8) converges to  $\boldsymbol{\nu}$  with respect to the adelic KR-topology.*

*Proof.* Put  $\boldsymbol{\nu} = (\nu_v)_v$  and let  $\varepsilon > 0$  be given. Let  $S$  be a finite nonempty subset of  $\mathfrak{M}_{\mathbb{K}}$  such that  $\nu_v = \delta_0$  for all  $v \notin S$ , and put

$$\varepsilon' = \frac{\varepsilon}{6 \sum_{v \in S} n_v} \quad \text{and} \quad d' = \frac{2}{\varepsilon'} \max_{v \in S} \int \|u\| \, d\nu_v.$$

By [Vil09, Theorem 6.18], for each  $v \in S$  we can approach  $\nu_v$  with respect to the KR-distance, by a probability measure with finite support. Therefore we can find  $d \geq \max\{2, d'\}$  and for each  $v \in S$  a sequence of points  $u_{v,1}, \dots, u_{v,d-1} \in N_{\mathbb{R}}$  such that the probability measure  $\nu_v'' = \frac{1}{d-1} \sum_{i=1}^{d-1} \delta_{u_{v,i}}$  satisfies  $W(\nu_v, \nu_v'') < \varepsilon'$ . We deduce from [Vil09, Formula (6.3)] the inequalities

$$\left| \int \|u\| \, d\nu_v - \int \|u\| \, d\nu_v'' \right| \leq \varepsilon' \quad \text{and} \quad \|\mathbb{E}[\nu_v] - \mathbb{E}[\nu_v'']\| \leq \varepsilon'.$$

Defining  $u_{v,d} := d\mathbb{E}[\nu_v] - (d-1)\mathbb{E}[\nu_v'']$ , we verify  $\|u_{v,d}\| \leq \|\mathbb{E}[\nu_v]\| + (d-1)\varepsilon'$ . Thus, setting  $\nu_v' = \frac{1}{d} \sum_{i=1}^d \delta_{u_{v,i}}$  and using Jensen's inequality and [Vil09, Formula (6.3)] again, we get

$$\begin{aligned} W(\nu_v'', \nu_v') &\leq \frac{1}{d(d-1)} \sum_{i=1}^{d-1} \|u_{v,i}\| + \frac{1}{d} \|u_{v,d}\| \\ &\leq \frac{1}{d} \left( \int \|u\| \, d\nu_v'' + \int \|u\| \, d\nu_v + (d-1)\varepsilon' \right) \leq \frac{2}{d} \int \|u\| \, d\nu_v + \varepsilon' \leq 2\varepsilon'. \end{aligned}$$

We then easily check  $\mathbb{E}[\nu_v'] = \mathbb{E}[\nu_v]$  and

$$W(\nu_v, \nu_v') \leq W(\nu_v, \nu_v'') + W(\nu_v'', \nu_v') < 3\varepsilon'.$$

Set also  $\nu_v' = \delta_0$  for  $v \notin S$ . Then  $\boldsymbol{\nu}' = (\nu_v')_v \in \mathcal{H}_{\mathbb{K}}$  and  $W_{\mathbb{K}}(\boldsymbol{\nu}, \boldsymbol{\nu}') < \frac{\varepsilon}{2}$ .

Let  $\mathbb{F}/\mathbb{K}$  be a finite extension of degree  $d$  such that all places in  $S$  split completely, as given by [BPS15, Lemma 2.2]. For each  $v \in S$  and  $w \in \mathfrak{M}_{\mathbb{F}}$  such that  $w \mid v$ , we have  $n_w = n_v/d$ . We enumerate the places above a given place  $v \in S$  as  $w(v, j)$ ,  $j = 1, \dots, d$ .

Let  $H_{\mathbb{F}} \subset \bigoplus_{w \in \mathfrak{M}_{\mathbb{F}}} N_{\mathbb{R}}$  be the subspace defined by the equation  $\sum_w n_w u_w = 0$ . For each  $v \in \mathfrak{M}_{\mathbb{K}}$  consider the element  $\mathbf{u} \in H_{\mathbb{F}}$  given, for  $w \in \mathfrak{M}_{\mathbb{F}}$ , by

$$u_w = \begin{cases} u_{v,j} & \text{for } v \in S \text{ and } w = w(v, j) \text{ with } 1 \leq j \leq d, \\ 0 & \text{for } v \notin S \text{ and } w \mid v. \end{cases}$$

Consider the map  $\text{val}_{\mathbb{F}}: \mathbb{T}(\mathbb{F}) \rightarrow \bigoplus_{w \in \mathfrak{M}_{\mathbb{F}}} N_{\mathbb{R}}$  defined by  $\text{val}_{\mathbb{F}} = (\text{val}_w)_{w \in \mathfrak{M}_{\mathbb{F}}}$ . This is a group homomorphism and so it can be extended to a map

$$\text{val}_{\mathbb{F}}: \mathbb{T}(\mathbb{F}) \otimes \mathbb{Q} \longrightarrow \bigoplus_{w \in \mathfrak{M}_{\mathbb{F}}} N_{\mathbb{R}}.$$

By the product formula, the image of this map lies in the hyperplane  $H_{\mathbb{F}}$  and, by [BPS15, Lemma 2.3], it is dense with respect to the  $L^1$ -topology on  $H_{\mathbb{F}}$ . For  $\alpha \in \mathbb{T}(\mathbb{F})$  and  $r \in \mathbb{Q}$ , we have

$$\begin{aligned} \|\mathbf{u} - \text{val}_{\mathbb{F}}(\alpha^r)\|_{L^1} &= \sum_{v \in S} \frac{n_v}{d} \sum_{j=1}^d \|u_{v,j} - \text{val}_{w(v,j)}(\alpha^r)\| + \sum_{v \notin S} \|\text{val}_v(\alpha^r)\| \\ &= \sum_v n_v \int \|u - u'\| d\lambda_v(u, u') \end{aligned} \quad (4.17)$$

for the probability measure  $\lambda_v$  on  $N_{\mathbb{R}} \times N_{\mathbb{R}}$  given by

$$\lambda_v = \begin{cases} \frac{1}{d} \sum_{j=1}^d \delta_{(u_{v,j}, \text{val}_{w(v,j)}(\alpha^r))} & \text{if } v \in S, \\ \delta_{(0, \text{val}_v(\alpha^r))} & \text{if } v \notin S. \end{cases}$$

This measure has marginals  $\nu'_v$  and  $\nu_{p,v}$  for any  $p = \omega \cdot \alpha^r$  with  $\omega$  a torsion point in  $\mathbb{T}(\overline{\mathbb{K}})$ , thus  $W(\nu'_v, \nu_{p,v}) \leq \int \|u - u'\| d\lambda_v(u, u')$  for every  $v$ , and the quantity in (4.17) is an upper bound for the adelic KR-distance  $W_{\mathbb{K}}(\boldsymbol{\nu}', \boldsymbol{\nu}_p)$ . It follows that we can choose  $\alpha$  and  $r$  such that  $W_{\mathbb{K}}(\boldsymbol{\nu}', \boldsymbol{\nu}_p) < \varepsilon/2$  and thus  $W_{\mathbb{K}}(\boldsymbol{\nu}, \boldsymbol{\nu}_p) < \varepsilon$ .

Since the orbit of  $\alpha^r$  under the action of the group of torsion points of  $\mathbb{T}(\overline{\mathbb{K}})$  is Zariski dense, we have shown that, given  $\varepsilon > 0$  and a nonempty open subset  $U \subset X$ , we can choose  $p \in U(\overline{\mathbb{K}})$  satisfying

$$W_{\mathbb{K}}(\boldsymbol{\nu}, \boldsymbol{\nu}_p) < \varepsilon.$$

As in the proof of Proposition 2.5, let  $I$  be the set of closed subvarieties of pure codimension 1 in  $X$  ordered by inclusion. For each  $Y \in I$  choose a point  $p_Y \in (X \setminus Y)(\overline{\mathbb{K}})$  such that

$$W_{\mathbb{K}}(\boldsymbol{\nu}, \boldsymbol{\nu}_{p_Y}) < \frac{1}{c(Y)}$$

with  $c(Y)$  the number of components of  $Y$ . Thus, the net of algebraic points  $(p_Y)_{Y \in I}$  is generic and the net of probability measures  $(\boldsymbol{\nu}_{p_Y})_{Y \in I}$  converges to  $\boldsymbol{\nu}$  in the KR-topology, proving the result.  $\square$

*Proof of Theorem 4.7.* Let  $\boldsymbol{\nu} = (\nu_v)_v$  be a centered adelic measure on  $N_{\mathbb{R}}$  such that each measure  $\nu_v$  satisfies the condition (4.5). By Lemma 4.8, it satisfies

$$\eta_{\overline{D}}(\boldsymbol{\nu}) = \mu_{\overline{D}}^{\text{ess}}(X).$$

Proposition 4.11 implies that there is a generic net  $(p_l)_{l \in I}$  of points in  $\mathbb{T}(\overline{\mathbb{K}}) = X_0(\overline{\mathbb{K}})$  such that  $(\boldsymbol{\nu}_{p_l})_{l \in I}$  converges to  $\boldsymbol{\nu}$  with respect to the distance  $W_{\mathbb{K}}$ . On the other hand, by Lemma 4.8 we also have

$$\lim_l h_{\overline{D}}(p_l) = \lim_l \eta_{\overline{D}}(\boldsymbol{\nu}_{p_l}) = \eta_{\overline{D}}(\boldsymbol{\nu}) = \mu_{\overline{D}}^{\text{ess}}(X),$$

and so the net  $(p_l)_{l \in I}$  is  $\overline{D}$ -small.  $\square$

**Corollary 4.12.** *Let  $v \in \mathfrak{M}_{\mathbb{K}}$ . For every measure  $\nu_v \in \mathcal{E}$  with  $\text{supp}(\nu_v) \subset F_v$  and  $E[\nu_v] \in B_v$ , there is a generic  $\overline{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$  such*

that the net of measures  $(\nu_{p_l, v})_{l \in I}$  converges to  $\nu_v$  with respect to the Kantorovich-Rubinstein distance. In particular,  $(\nu_{p_l, v})_{l \in I}$  also converges to  $\nu_v$  in the weak-\* topology with respect to  $\mathcal{C}_b(N_{\mathbb{R}})$ .

*Proof.* Let  $x$  be a point where  $\vartheta_{\overline{D}}$  attains its maximum. Since

$$\mathbb{E}[\nu_v] \in B_v \subset -\partial \left( \sum_{w \in \mathfrak{M}_{\mathbb{K}} \setminus \{v\}} \frac{n_w}{n_v} \vartheta_{\overline{D}, w} \right) (x) = - \sum_{w \in \mathfrak{M}_{\mathbb{K}} \setminus \{v\}} \frac{n_w}{n_v} \partial \vartheta_{\overline{D}, w} (x),$$

we can find  $u_w \in \partial \vartheta_{\overline{D}, w}$  for each  $w \neq v$  such that

$$u_v := \mathbb{E}[\nu_v] = - \sum_{w \in \mathfrak{M}_{\mathbb{K}} \setminus \{v\}} \frac{n_w}{n_v} u_w.$$

In particular, for all  $w \in \mathfrak{M}_{\mathbb{K}}$  one has

$$u_w = - \sum_{w' \in \mathfrak{M}_{\mathbb{K}} \setminus \{w\}} \frac{n_{w'}}{n_w} u_{w'} \in \partial \vartheta_{\overline{D}, w} (x) \cap \left( -\partial \left( \sum_{w' \in \mathfrak{M}_{\mathbb{K}} \setminus \{w\}} \frac{n_{w'}}{n_w} \vartheta_{\overline{D}, w'} \right) (x) \right) = B_w.$$

Furthermore, we have  $u_w = 0$  for all but a finite number of places  $w$  in  $\mathfrak{M}_{\mathbb{K}}$ . Put  $\nu_w = \delta_{u_w}$  for each  $w \neq v$ . The statement then follows from Theorem 4.7 applied to the centered adelic measure  $\nu = (\nu_w)_{w \in \mathfrak{M}_{\mathbb{K}}}$ .  $\square$

Combining Theorems 4.3 and 4.7, we can obtain a criterion for when the direct image under the valuation map of the Galois orbits of a small net converges in the sense of measures. We show that in this case, the limit measure is concentrated in a single point.

**Corollary 4.13.** *Let  $v \in \mathfrak{M}_{\mathbb{K}}$ . The following conditions are equivalent:*

- (1) *for every  $\overline{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$ , the net of measures  $(\nu_{p_l, v})_{l \in I}$  converges in the weak-\* topology with respect to  $\mathcal{C}_b(N_{\mathbb{R}})$ ;*
- (2) *for every generic  $\overline{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$ , the net of measures  $(\nu_{p_l, v})_{l \in I}$  converges in the weak-\* topology with respect to  $\mathcal{C}_c(N_{\mathbb{R}})$ , the space of continuous functions on  $N_{\mathbb{R}}$  with compact support;*
- (3) *the face  $F_v$  contains only one point.*

*When these equivalent conditions hold, the limit measures in (1) and (2) coincide with the Dirac measure at the unique point of  $F_v$ .*

*Proof.* It is clear that (1) implies (2), and Theorem 4.3 shows that (3) implies (1). Now suppose that the face  $F_v$  has more than one point. Since  $F_v$  is the minimal face containing  $B_v$ , we can find distinct points  $u_0, u_1, u_2 \in F_v$  such that

$$u_0 = \frac{u_1 + u_2}{2} \in B_v.$$

The probability measures  $\delta_{u_0}$  and  $\frac{1}{2}\delta_{u_1} + \frac{1}{2}\delta_{u_2}$  satisfy the conditions (4.5). By Corollary 4.12, we can find generic  $\overline{D}$ -small nets  $(p_l)_{l \in I}$  and  $(q_l)_{l \in I}$  such that the nets of measures  $(\nu_{p_l, v})_{l \in I}$  and  $(\nu_{q_l, v})_{l \in I}$  respectively converge to

$$\delta_{u_0} \quad \text{and} \quad \frac{1}{2}\delta_{u_1} + \frac{1}{2}\delta_{u_2}$$

in the KR-topology, and hence in the weak-\* topology with respect to  $\mathcal{C}_c(N_{\mathbb{R}})$ . Combining these nets, we can obtain a net that does not converge in this weak-\* topology. Hence the condition (2) implies the condition (3).

The last statement follows from Theorem 4.3.  $\square$

When any of the equivalent conditions of Corollary 4.13 holds we say that the metrized divisor  $\overline{D}$  satisfies the *modulus concentration property* at the place  $v$ . Thus Corollary 4.13 gives us a criterion for the modulus concentration property

at a place. We next give a criterion for the modulus concentration property at all places simultaneously, which can be directly read from the roof function. Before giving it, we need some preliminary results and a definition.

**Definition 4.14.** A semipositive toric metrized  $\mathbb{R}$ -divisor  $\overline{D}$  with  $D$  big is called *monocritical* if the minimum of  $\eta_{\overline{D}}$  in  $\mathcal{H}_{\mathbb{K}}$  is attained at a unique point. If this is the case, by Corollary 4.10, the minimum is attained at a point of  $H_{\mathbb{K}}$ . This point is called *the critical point of  $\overline{D}$* .

**Example 4.15.** Let  $\overline{D}^{\text{can}}$  be a nef and big toric  $\mathbb{R}$ -divisor equipped with the canonical metric as in Example 4.1. Then all its local roof functions are zero. Taking a point  $x$  in the interior of the polytope, we have  $\partial\vartheta_{\overline{D},v}(x) = \{0\}$  for every  $v$ . Hence  $F_v = \{0\}$  for every  $v$  and  $\overline{D}$  is monocritical with critical point  $\mathbf{0} \in H_{\mathbb{K}}$ .

Recall that  $\Delta_{D,\max}$  denotes the convex set of points of  $\Delta_D$  where  $\vartheta_{\overline{D}}$  attains its maximum.

**Proposition 4.16.** *The following conditions are equivalent:*

- (1) *the metrized  $\mathbb{R}$ -divisor  $\overline{D}$  is monocritical;*
- (2) *for every point  $x \in \Delta_{D,\max}$ , the set*

$$H_{\mathbb{K}} \cap \prod_{v \in \mathfrak{M}_{\mathbb{K}}} \partial\vartheta_{\overline{D},v}(x) \quad (4.18)$$

*contains a unique element  $\mathbf{u} = (u_v)_v \in H_{\mathbb{K}}$  and, for  $v \in \mathfrak{M}_{\mathbb{K}}$ , the point  $u_v$  is a vertex of  $\partial\vartheta_{\overline{D},v}(x)$ ;*

- (3) *for every point  $x \in \Delta_{D,\max}$ , the point  $0$  is a vertex of  $\partial\vartheta_{\overline{D}}(x)$ ;*
- (4) *there exists a point  $x \in \Delta_{D,\max}$  such that  $0$  is a vertex of  $\partial\vartheta_{\overline{D}}(x)$ ;*
- (5) *for all  $v \in \mathfrak{M}_{\mathbb{K}}$ , the set  $F_v$  contains only one point.*

*When these equivalent conditions hold,  $F_v = \{u_v\}$  for every  $v$  and  $\mathbf{u}$  is the critical point of  $\overline{D}$ .*

*Proof.* We prove first that (1) implies (2). Assume that  $\overline{D}$  is monocritical. Let  $\mathbf{u} = (u_v)_v$  belong to the set (4.18). Then for every  $v \in \mathfrak{M}_{\mathbb{K}}$  we have

$$u_v \in \partial\vartheta_{\overline{D},v}(x) \cap \left( -\partial \left( \sum_{w \in \mathfrak{M}_{\mathbb{K}}, w \neq v} \frac{n_w}{n_v} \vartheta_{\overline{D},w} \right) (x) \right).$$

So the measure  $\nu = (\delta_{u_v})_v$  belongs to  $\mathcal{H}_{\mathbb{K}}$  and satisfies  $\text{supp}(\delta_{u_v}) \subset B_v$  for each  $v$ . In particular,  $\text{supp}(\delta_{u_v}) \subset F_v$  and  $E[\delta_{u_v}] \in B_v$ . Thus by Lemma 4.8

$$\eta_{\overline{D}}(\mathbf{u}) = \min_{\nu' \in \mathcal{H}_{\mathbb{K}}} \eta_{\overline{D}}(\nu').$$

Since  $\overline{D}$  is monocritical, this shows that the set (4.18) is reduced to the unique critical point of  $\overline{D}$ .

Assume now that the set (4.18) contains a single point  $\mathbf{u} = (u_v)_v \in H_{\mathbb{K}}$  and there is a place  $v_0 \in \mathfrak{M}_{\mathbb{K}}$  such that  $u_{v_0}$  is not a vertex of  $\partial\vartheta_{\overline{D},v_0}(x)$ . Then we can find two points  $u_{v_0,1}, u_{v_0,2} \in \partial\vartheta_{\overline{D},v_0}(x)$  such that

$$u_{v_0} = \frac{u_{v_0,1} + u_{v_0,2}}{2}.$$

We consider the measure  $\nu_1 = (\delta_{u_v})_v$  and the measure  $\nu_2 = (\nu_v)_v$  defined by

$$\nu_v = \begin{cases} \delta_{u_v} & \text{if } v \neq v_0, \\ \frac{\delta_{u_{v_0,1}} + \delta_{u_{v_0,2}}}{2} & \text{if } v = v_0. \end{cases}$$

Then  $\nu_2$  is in (4.18) and, again by Lemma 4.8, we have that

$$\eta_{\overline{D}}(\nu_1) = \eta_{\overline{D}}(\nu_2) = \min_{\nu \in \mathcal{H}_{\mathbb{K}}} \eta_{\overline{D}}(\nu)$$

contradicting the hypothesis that  $\overline{D}$  is monocritical, and completing the proof of (2).

Assume that (2) is true and fix  $x \in \Delta_{D, \max}$ . Let  $S \subset \mathfrak{M}_{\mathbb{K}}$  be the finite set of places where  $u_v \neq 0$  or  $\vartheta_{\overline{D}, v}$  is not identically zero. We have that

$$\partial \vartheta_{\overline{D}}(x) = \sum_{v \in S} n_v \partial \vartheta_{\overline{D}, v}(x).$$

Moreover, (2) implies that the equation

$$0 = \sum_{v \in S} n_v a_v \quad \text{with } a_v \in \partial \vartheta_{\overline{D}, v}(x)$$

has a unique solution  $a_v = u_v$  and this solution satisfies that  $a_v$  is a vertex of  $\partial \vartheta_{\overline{D}, v}(x)$ . Therefore, by Lemma 3.15 we deduce that 0 is a vertex of  $\partial \vartheta_{\overline{D}}(x)$ . Hence (2) implies (3).

Since  $\Delta_{D, \max}$  is nonempty, (3) implies (4).

Assume now that (4) is true. For each  $v$ , let  $g_{1, v}$  and  $g_{2, v}$  be the continuous concave functions on  $\Delta_D$  in Notation 4.2. Since  $\vartheta_{\overline{D}} = n_v g_{1, v} + n_v g_{2, v}$ ,

$$\partial \vartheta_{\overline{D}}(x) = n_v \partial g_{1, v}(x) + n_v \partial g_{2, v}(x).$$

Lemma 3.15 and the definition of the set  $B_v$  imply that this set contains one single point  $u_v$ , and that this point is a vertex of both  $\partial g_{1, v}(x)$  and of  $-\partial g_{2, v}(x)$ . Hence  $B_v$  is already a face of  $\partial g_{1, v}(x)$ . Thus  $F_v = B_v = \{u_v\}$  and so (4) implies (5).

By Lemma 4.8 it is clear that (5) implies (1) finishing the proof of the equivalence.

Assume now that  $\overline{D}$  is monocritical. Since by Lemma 4.8 the point  $\mathbf{u}$  in (2) satisfies that  $\eta_{\overline{D}}(\mathbf{u}) = \min_{\nu \in \mathcal{H}_{\mathbb{K}}} \eta_{\overline{D}}(\nu)$ , it is the critical point. Following the proof of the equivalence we deduce that  $F_v = \{u_v\}$  proving the last statement.  $\square$

For a given toric metrized  $\mathbb{R}$ -divisor, the condition of being monocritical and its critical point behave well with respect to scalar extensions. The following result follows from the compatibility of toric metrics with scalar extensions in [BPS14, Proposition 4.3.8].

**Proposition 4.17.** *Let  $X$  and  $\overline{D}$  as before. Let  $\mathbb{F} \subset \overline{\mathbb{K}}$  be a finite extension of  $\mathbb{K}$  and write  $\overline{D}_{\mathbb{F}}$  for the toric metrized  $\mathbb{R}$ -divisor on  $X_{\mathbb{F}}$  obtained by scalar extension. If  $\overline{D}$  is monocritical with critical point  $(u_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ , then  $\overline{D}_{\mathbb{F}}$  is also monocritical and its critical point  $(u_w)_{w \in \mathfrak{M}_{\mathbb{F}}}$  is given by  $u_w = u_v$  for all  $v \in \mathfrak{M}_{\mathbb{K}}$  and  $w$  over  $v$ .*

We now give the criterion for modulus concentration at every place.

**Theorem 4.18.** *Let  $X$  and  $\overline{D}$  be as before. The following conditions are equivalent:*

- (1) *for every  $\overline{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$  and every place  $v \in \mathfrak{M}_{\mathbb{K}}$ , the net of measures  $(\nu_{p_l, v})_{l \in I}$  converges.*
- (2) *the metrized  $\mathbb{R}$ -divisor  $\overline{D}$  is monocritical;*

*When these equivalent conditions hold,*

$$\lim_{l \in I} \nu_{p_l, v} = \delta_{u_v},$$

*where  $(u_v)_v$  is the critical point of  $\overline{D}$ .*

*Proof.* The theorem follows directly from Corollary 4.13 and Proposition 4.16.  $\square$

When there is modulus concentration for every place, we can show that the convergence holds not only in the weak-\* topology with respect to  $\mathcal{C}_b(N_{\mathbb{R}})$  but even in the stronger adelic KR-topology.

**Theorem 4.19.** *Let  $X$  and  $\overline{D}$  be as before. Assume that  $\overline{D}$  is monocritical. Let  $\mathbf{u} = (u_v)_v$  be the critical point of  $\overline{D}$  and set  $\delta_{\mathbf{u}} = (\delta_{u_v})_v \in \mathcal{H}_{\mathbb{K}}$ . Then, for every  $\overline{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$ , the net of centered adelic measures  $(\nu_{p_l})_{l \in I}$  converges to  $\delta_{\mathbf{u}}$  in the adelic KR-topology. In particular, for every  $v \in \mathfrak{M}_{\mathbb{K}}$ , the net of measures  $(\nu_{p_l, v})_{l \in I}$  converges to  $\delta_{u_v}$  in the KR-topology.*

*Proof.* For each  $v \in \mathfrak{M}_{\mathbb{K}}$ , let  $f_v: N_{\mathbb{R}} \rightarrow \mathbb{R}$  be the function given by

$$f_v(u) = \psi_{\overline{D}, v}(u) - \Psi_D(u - u_v).$$

This is an adelic family of bounded continuous functions on  $N_{\mathbb{R}}$  with  $f_v = 0$  for all but a finite number of  $v$ . Consider then the function  $\eta': \mathcal{H}_{\mathbb{K}} \rightarrow \mathbb{R}$  given by

$$\eta'(\nu) = \eta_{\overline{D}}(\nu) + \sum_v n_v \int f_v d\nu_v = - \sum_v n_v \int \Psi_D(u - u_v) d\nu_v.$$

Since the net  $(p_l)_{l \in I}$  is  $\overline{D}$ -small,

$$\lim_l \eta_{\overline{D}}(\nu_{p_l}) = \lim_l h_{\overline{D}}(p_l) = \mu_{\overline{D}}^{\text{ess}}(X).$$

By Theorem 4.18, for every place  $v \in \mathfrak{M}_{\mathbb{K}}$  the net of measures  $(\nu_{p_l, v})_{l \in I}$  converges to  $\delta_{u_v}$ , so that  $\lim_l \int f_v d\nu_{p_l, v} = \int f_v d\delta_{u_v} = \psi_{\overline{D}, v}(u_v)$ . Since  $\mathbf{u} = (u_v)_v$  is the critical point of  $\overline{D}$ , using Corollary 4.10 we get

$$\lim_l \eta'(\nu_{p_l}) = \mu_{\overline{D}}^{\text{ess}}(X) + \sum_v n_v \psi_{\overline{D}, v}(u_v) = 0. \quad (4.19)$$

Choose a point  $x$  in the interior of  $\Delta_D$ . Then there is a constant  $c > 0$  such that, for all  $u \in N_{\mathbb{R}}$ ,

$$\|u\| \leq -c(\Psi_D - x)(u).$$

It follows from the definition of the Kantorovich-Rubinstein distance that, for each  $v \in \mathfrak{M}_{\mathbb{K}}$ ,

$$W(\nu_{p_l, v}, \delta_{u_v}) \leq \int \|u - u_v\| d\nu_{p_l, v}(u).$$

Hence

$$\begin{aligned} W_{\mathbb{K}}(\nu_{p_l}, \delta_{\mathbf{u}}) &\leq \sum_v n_v \int \|u - u_v\| d\nu_{p_l, v}(u) \\ &\leq -c \sum_v n_v \int (\Psi_D - x)(u - u_v) d\nu_{p_l, v}(u) = c\eta'(\nu_{p_l}), \end{aligned}$$

where the last equality follows from the facts that  $\mathbf{u}$  belongs to  $H_{\mathbb{K}}$  and that  $\nu_{p_l}$  is a centered adelic measure on  $N_{\mathbb{R}}$ , thanks to the product formula in Proposition 2.1(2). By (4.19), this distance converges to 0, completing the proof.  $\square$

## 5. EQUIDISTRIBUTION OF GALOIS ORBITS AND THE BOGOMOLOV PROPERTY

We turn to the study of the limit measures of Galois orbits of  $\overline{D}$ -small nets of algebraic points in toric varieties. In this section, we denote by  $X$  a proper toric variety over a global field  $\mathbb{K}$  and  $\overline{D}$  a toric metrized  $\mathbb{R}$ -divisor on  $X$  with  $D$  big. For  $v \in \mathfrak{M}_{\mathbb{K}}$ , recall that  $\text{val}_v: \mathbb{T}_v^{\text{an}} \rightarrow N_{\mathbb{R}}$  denotes the valuation map, defined in (4.1).

We first describe the limit measures in the monocritical case.

**Definition 5.1.** Given  $v \in \mathfrak{M}_{\mathbb{K}}$  and  $u \in N_{\mathbb{R}}$ , the probability measure  $\lambda_{\mathbb{S}_v, u}$  on  $X_v^{\text{an}}$  is defined as follows.



- (1) When  $v$  is Archimedean, note that  $\text{val}_v^{-1}(u) = \mathbb{S}_v \cdot p$  for any point  $p \in \text{val}_v^{-1}(u)$  and where  $\mathbb{S}_v = \text{val}_v^{-1}(0) \simeq (S^1)^n$  is the compact torus of  $\mathbb{T}_v^{\text{an}}$ . In this case,  $\lambda_{\mathbb{S}_v, u}$  is the direct image under the translation by  $p$  of the Haar probability measure of  $\mathbb{S}_v$ .
- (2) When  $v$  is non-Archimedean, consider the multiplicative seminorm on the group algebra  $\mathbb{C}_v[M] \simeq \mathbb{C}_v[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$  that, to a Laurent polynomial  $\sum_{m \in M} \alpha_m \chi^m$ , assigns the value  $\max_m (|\alpha_m|_v e^{-\langle m, u \rangle})$ . This seminorm gives a point, denoted by  $\theta(u)$ , in the Berkovich space  $X_v^{\text{an}}$ . The point  $\theta(u)$  lies in the preimage  $\text{val}_v^{-1}(u)$ . We then set  $\lambda_{\mathbb{S}_v, u} = \delta_{\theta(u)}$ , the Dirac measure at this point.

The following result corresponds to Theorem 1.1 in the introduction, and shows that modulus concentration at every place implies the equidistribution property at every place. Due to the existing equidistribution theorems in the literature, we restrict its statement to divisors (rather than  $\mathbb{R}$ -divisors).

**Theorem 5.2.** *Let  $X$  be a proper toric variety over  $\mathbb{K}$  and  $\overline{D}$  a semipositive toric metrized divisor on  $X$  with  $D$  big. The following conditions are equivalent:*

- (1) *for every generic  $\overline{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$  and every place  $v \in \mathfrak{M}_{\mathbb{K}}$ , the net of probability measures  $(\mu_{p_l, v})_{l \in I}$  on  $X_v^{\text{an}}$  converges;*
- (2) *the metrized divisor  $\overline{D}$  is monocritical.*

*When these equivalent conditions hold, the limit measure in (1) is  $\lambda_{\mathbb{S}_v, u_v}$ , with  $u_v \in N_{\mathbb{R}}$  the  $v$ -adic component of the critical point of  $\overline{D}$ .*

The proof of Theorem 5.2 is done by reduction to the quasi-canonical case. The following is the characterization of quasi-canonical toric metrized  $\mathbb{R}$ -divisors in [BPS15].

**Proposition 5.3.** *Let  $X$  be a proper toric variety over  $\mathbb{K}$  and  $\overline{D}$  a semipositive toric metrized  $\mathbb{R}$ -divisor on  $X$  with  $D$  big. The following conditions are equivalent:*

- (1)  *$\overline{D}$  is quasi-canonical (Definition 2.7);*
- (2)  *$\vartheta_{\overline{D}}$  is constant;*
- (3) *there are  $\mathbf{u} = (u_v)_v \in H_{\mathbb{K}}$  and  $(\gamma_v)_v \in \bigoplus_{v \in \mathfrak{M}_{\mathbb{K}}} \mathbb{R}$  such that*

$$\psi_{\overline{D}, v}(u) = \Psi_D(u - u_v) - \gamma_v$$

*for all  $v \in \mathfrak{M}_{\mathbb{K}}$  and  $u \in N_{\mathbb{R}}$ .*

*Proof.* The equivalence of (1) and (3) is given by [BPS15, Corollary 4.7]. The equivalence of (1) and (2) is given in the course of the proof of [BPS15, Proposition 4.6], recalling that  $\text{vol}(D) = \text{deg}_D(X)$  and noting that, since by assumption  $\overline{D}$  is semipositive,  $\widehat{\text{vol}}_{\chi}(\overline{D}) = h_{\overline{D}}(X)$ .  $\square$

The following result gives the key step in the proof of Theorem 5.2.

**Proposition 5.4.** *Let  $X$  be a proper toric variety over  $\mathbb{K}$  and  $\overline{D}$  a monocritical metrized  $\mathbb{R}$ -divisor on  $X$  with critical point  $\mathbf{u} = (u_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ . Let  $\overline{D}'$  be the toric metrized  $\mathbb{R}$ -divisor over  $D$  corresponding to the family of concave functions  $\psi_{\overline{D}', v}: N_{\mathbb{R}} \rightarrow \mathbb{R}$ ,  $v \in \mathfrak{M}_{\mathbb{K}}$ , given by*

$$\psi_{\overline{D}', v}(u) = \Psi_D(u - u_v). \quad (5.1)$$

*Then  $\overline{D}'$  is quasi-canonical and every  $\overline{D}$ -small net of algebraic points of  $X_0$  is also  $\overline{D}'$ -small.*

*Proof.* The fact that  $\overline{D}'$  is quasi-canonical is given by Proposition 5.3.

Let  $(p_l)_{l \in I}$  be a  $\overline{D}$ -small net of algebraic points of  $X_0$ . By Theorem 4.19, the net of centered adelic measures  $(\nu_{p_l})_{l \in I}$  converges to  $\delta_{\mathbf{u}} = (\delta_{u_v})_v$  with respect to the

adelic KR-distance. By Lemma 4.8, the function  $\eta_{\overline{D}'}$  is continuous with respect to this distance. Using (4.9), we deduce that

$$\lim_l h_{\overline{D}'}(p_l) = \lim_l \eta_{\overline{D}'}(\nu_{p_l}) = \eta_{\overline{D}'}(\delta_u) = 0.$$

On the other hand,  $\vartheta_{\overline{D}',v} = u_v$  for each  $v$ . Since the critical point  $u$  lies in the subspace  $H_{\mathbb{K}}$ , we have that  $\vartheta_{\overline{D}'} = \sum_v n_v u_v = 0$ . Hence,

$$\mu_{\overline{D}'}^{\text{ess}}(X) = \max_{x \in \Delta_D} \vartheta_{\overline{D}'}(x) = 0.$$

Thus  $(p_l)_{l \in I}$  is  $\overline{D}'$ -small, as stated.  $\square$

*Proof of Theorem 5.2.* Suppose that the condition (1) holds. Given a generic  $\overline{D}$ -small net  $(p_l)_{l \in I}$  of algebraic points of  $X_0$  and  $v \in \mathfrak{M}_{\mathbb{K}}$ , the net of measures  $(\mu_{p_l,v})_{l \in I}$  converges weakly with respect to the space  $\mathcal{C}(X_v^{\text{an}})$ . Hence, the net of direct images  $(\nu_{p_l,v})_{l \in I}$  converges weakly with respect to the space  $\mathcal{C}_c(N_{\mathbb{R}})$ . By Corollary 4.13, for each  $v$ , the face  $F_v$  contains only one point. Proposition 4.16 then implies that  $\overline{D}$  is monocritical, giving the condition (2).

Now suppose that the condition (2) holds. Since  $\overline{D}$  is monocritical, the polytope  $\Delta_D$  has nonempty interior. Let  $Y$  be the toric variety associated to the normal fan of  $\Delta_D$  and  $E$  the divisor on  $Y$  associated to the virtual support function  $\Psi_D$ , see for example [BPS14, Theorem 3.3.3]. By construction  $E$  is ample and  $(Y, E)$  is the polarized toric variety associated to the polytope  $\Delta_D$ , see for example [BPS14, Theorem 3.4.6 and Remark 3.4.7]. By the characterization of semipositive toric metrics in [BPS14, Theorem 4.8.1], the metric in  $\overline{D}$  induces a semipositive toric metric on  $E$ , and we denote by  $\overline{E}$  the corresponding toric metrized divisor. We have that  $\psi_{\overline{E},v} = \psi_{\overline{D},v}$  for all  $v$ , and so  $\overline{E}$  is also monocritical with the same critical point as  $\overline{D}$ .

Let

$$\overline{E}' = (E, \|\cdot\|'_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$$

be the ample divisor  $E$  on  $Y$  equipped with the quasi-canonical toric metric given by Proposition 5.4, with  $\overline{D}$  replaced by  $\overline{E}$ . Let  $(p_l)_{l \in I}$  be a generic  $\overline{D}$ -small net of algebraic points of  $X_0 = \mathbb{T} = Y_0$ . It is also a generic  $\overline{E}$ -small net of algebraic points of  $Y_0$ . By Proposition 5.4 with  $\overline{D}$  replaced by  $\overline{E}$ , it is also  $\overline{E}'$ -small.

By Theorem 2.11, for each place  $v$  the net  $(\mu_{p_l,v})_{l \in I}$  converges to the normalized Monge-Ampère measure  $\mu_v = \frac{1}{\deg_E(Y)} c_1(E, \|\cdot\|'_v)^{\wedge n}$  on  $Y_v^{\text{an}}$ . Consider the real Monge-Ampère measure  $\mathcal{M}(\psi_{\overline{E}',v})$  associated to the  $v$ -adic metric in  $\overline{E}'$  as in [BPS14, Definition 2.7.1]. By the explicit formula (5.1) and [BPS14, Example 2.7.5],

$$\mathcal{M}(\psi_{\overline{E}',v}) = \text{vol}_M(\Delta_D) \delta_{u_v} = \frac{\deg_E(Y)}{n!} \delta_{u_v}.$$

Then [BPS14, Theorem 4.8.11] implies that  $\mu_v = \lambda_{\mathbb{S}_v, u_v}$ . Therefore, the net of measures  $(\mu_{p_l,v})_{l \in I}$  on  $X_v^{\text{an}}$  converges to  $\lambda_{\mathbb{S}_v, u_v}$ , giving the condition (1) and the last statement in the theorem.  $\square$

**Example 5.5.** Let  $\overline{D}^{\text{can}}$  be a big and nef toric divisor on  $X$  equipped with the canonical metric. Following Example 4.15, this toric metrized divisor is monocritical with critical point  $\mathbf{0} \in H_{\mathbb{K}}$ . Hence, it satisfies the  $v$ -adic equidistribution property with limit measure  $\lambda_{\mathbb{S}_v, \mathbf{0}}$ , for every  $v \in \mathfrak{M}_{\mathbb{K}}$ .

In [Bil97], Bilu gave an equidistribution theorem for Galois orbits of sequences of points of small canonical height. This result is restricted to number fields and Archimedean places. However, and in contrast to the previous example, this result holds not just for generic, but for *strict* sequences of points, that is, sequences that eventually avoid any given proper torsion subvariety. This stronger version of

the equidistribution property was used in a crucial way in *loc. cit.* to prove the Bogomolov property for the canonical height.

Here we extend this version of the equidistribution property to monocritical metrized  $\mathbb{R}$ -divisors on toric varieties (Theorem 5.7) and deduce from it the Bogomolov property (Theorem 1.4 in the introduction, or Theorem 5.12 below). Our proofs are similar to Bilu's and use Fourier analysis. Hence, for the rest of the section we restrict to the case when  $\mathbb{K}$  is a number field and we only study the equidistribution over the Archimedean places. Following Remark 2.10, we restrict without loss of generality to sequences, instead of nets.

To formulate this extension, we have to modify slightly the notion of strict sequence. First we recall some standard terminology: a *subtorus* of  $\mathbb{T}$  is an algebraic subgroup of  $\mathbb{T}$  that is geometrically irreducible, a *translate of a subtorus* is a subvariety of  $\mathbb{T}_{\overline{\mathbb{K}}}$  that is the orbit of a point  $p \in \mathbb{T}(\overline{\mathbb{K}})$  by a subtorus, and a *torsion subvariety* is a translate of a subtorus by a torsion point of the group  $\mathbb{T}(\overline{\mathbb{K}}) \simeq (\overline{\mathbb{K}}^\times)^n$ .

**Definition 5.6.** A sequence  $(p_l)_{l \geq 1}$  of algebraic points of  $\mathbb{T}$  is *strict* if, for every translate of a subtorus  $U \subsetneq \mathbb{T}_{\overline{\mathbb{K}}}$ , there is  $l_0 \geq 1$  such that  $p_l \notin U(\overline{\mathbb{K}})$  for all  $l \geq l_0$ . Equivalently,  $(p_l)_{l \geq 1}$  is strict if, for every  $m \in M \setminus \{0\}$  and every point  $q \in X_0(\overline{\mathbb{K}})$ , there is  $l_0 \geq 1$  such that  $\chi^m(p_l) \neq \chi^m(q)$  for all  $l \geq l_0$ .

**Theorem 5.7.** *Let  $X$  be a proper toric variety over a number field  $\mathbb{K}$  and  $\overline{D}$  a monocritical metrized  $\mathbb{R}$ -divisor on  $X$ . Then, for every strict  $\overline{D}$ -small sequence  $(p_l)_{l \geq 1}$  of algebraic points of  $X_0$  and every Archimedean place  $v \in \mathfrak{M}_{\mathbb{K}}$ , the sequence  $(\mu_{p_l, v})_{l \geq 1}$  converges to the probability measure  $\lambda_{S_v, u_v}$ , with  $u_v \in N_{\mathbb{R}}$  the  $v$ -adic component of the critical point of  $\overline{D}$ .*

*Proof.* Let  $(p_l)_{l \geq 1}$  be a strict  $\overline{D}$ -small sequence of algebraic points of  $X_0$ . For each  $m \in M \setminus \{0\}$  consider the character

$$\chi^m: \mathbb{T} \longrightarrow \mathbb{G}_{m, \mathbb{K}}.$$

Since  $(p_l)_{l \geq 1}$  is strict, the sequence  $(\chi^m(p_l))_{l \geq 1}$  is generic.

We embed  $\mathbb{G}_{m, \mathbb{K}} \hookrightarrow \mathbb{P}_{\mathbb{K}}^1$  as the principal open subset. Let  $D_0 = \text{div}(x_0)$  be the divisor at infinity on  $\mathbb{P}_{\mathbb{K}}^1$ , equipped with the toric metric corresponding to the adelic family of functions  $\psi_{\overline{D}_0^m, v}: \mathbb{R} \rightarrow \mathbb{R}$  given by

$$\psi_{\overline{D}_0^m, v}(u) = \min(0, u - \langle m, u_v \rangle).$$

By Proposition 5.3, this metric is quasi-canonical. For each  $v \in \mathfrak{M}_{\mathbb{K}}$ , there is a commutative diagram

$$\begin{array}{ccc} \mathbb{T}_v^{\text{an}} & \xrightarrow{\chi^m} & \mathbb{G}_{m, v}^{\text{an}} \\ \text{val}_v \downarrow & & \downarrow \text{val}_v \\ N_{\mathbb{R}} & \xrightarrow{m} & \mathbb{R} \end{array}$$

The commutativity of this diagram implies that  $\nu_{\chi^m(p_l), v} = m_* \nu_{p_l, v}$ . By Theorem 4.19, the sequence  $(\nu_{p_l})_{l \geq 1}$  converges in the adelic KR-topology to the centered adelic measure  $(\delta_{u_v})_v$  on  $N_{\mathbb{R}}$ . Hence, the sequence  $(\nu_{\chi^m(p_l)})_{l \geq 1}$  converges in the adelic KR-topology to the centered adelic measure  $(\delta_{\langle m, u_v \rangle})_v$  on  $\mathbb{R}$ . By Lemma 4.8,

$$\lim_l \eta_{\overline{D}_0^m}(\nu_{\chi^m(p_l)}) = \eta_{\overline{D}_0^m}((\delta_{\langle m, u_v \rangle})_v) = \mu_{\overline{D}_0^m}^{\text{ess}}(\mathbb{P}_{\mathbb{K}}^1).$$

By the identity in (4.9),  $\eta_{\overline{D}_0^m}(\nu_{\chi^m(p_l)}) = h_{\overline{D}_0^m}(\chi^m(p_l))$ . Thus the sequence of points  $(\chi^m(p_l))_{l \geq 1}$  is  $\overline{D}_0^m$ -small.

Summarizing, the sequence  $(\chi^m(p_l))_{l \geq 1}$  of algebraic points of  $\mathbb{P}_{\mathbb{K}}^1$  is generic and small with respect to the quasi-canonical toric metrized divisor  $\overline{D}_0^m$ . Theorem 2.11

then implies that the sequence of measures  $(\mu_{\chi^m(p_l),v})_{l \geq 1}$  on the analytification  $\mathbb{P}_v^{\text{an}} \simeq \mathbb{P}^1(\mathbb{C})$  converges to  $\lambda_{\mathbb{S}_v, \langle m, u_v \rangle}$ .

Assume now that  $v$  is Archimedean. Since the space of probability measures on  $X(\mathbb{C})$  is sequentially compact, by restricting to a subsequence we can suppose without loss of generality that  $(\mu_{p_l, v})_{l \geq 1}$  converges to a measure  $\mu$ . Since the sequence of direct images  $((\text{val}_v)_* \mu_{p_l, v})_{l \geq 1}$  converges in the KR-topology to the Dirac measure on the point  $u_v \in N_{\mathbb{R}}$ , we deduce that

$$\text{supp}(\mu) \subset \text{val}_v^{-1}(u_v) = \mathbb{S}_v \cdot e^{-u_v}.$$

Let  $z$  be the standard affine coordinate of  $\mathbb{P}^1(\mathbb{C})$ . For each  $m \in M \setminus \{0\}$ , let  $z_m$  be a continuous function on  $\mathbb{P}^1(\mathbb{C})$  that agrees with  $z$  on a neighborhood of  $S^1 \cdot \chi^m(e^{-u_v})$ . Hence  $(\chi^m)^*(z_m)$  agrees with the character  $\chi^m$  on a neighborhood of  $\mathbb{S}_v \cdot e^{-u_v}$ . Then

$$\begin{aligned} \int \chi^m d\mu &= \int (\chi^m)^*(z_m) d\mu = \lim_l \int (\chi^m)^*(z_m) d\mu_{p_l, v} \\ &= \lim_l \int z_m d(\chi^m)_* \mu_{p_l, v} = \lim_l \int z_m d\mu_{\chi^m(p_l), v} \\ &= \int z_m d\lambda_{S^1, \langle m, u_v \rangle} = \int z d\lambda_{S^1, \langle m, u_v \rangle} = 0, \end{aligned}$$

where the last equality comes from Cauchy's formula. Hence  $\int \chi^m d\mu = 0$  for all  $m \in M \setminus \{0\}$ . By Fourier analysis, the only probability measure supported on  $\mathbb{S}_v \cdot e^{-u_v}$  satisfying this condition is  $\lambda_{\mathbb{S}_v, u_v}$ . Thus  $\mu = \lambda_{\mathbb{S}_v, u_v}$ , concluding the proof.  $\square$

**Remark 5.8.** Our notion of strict sequence is stronger than the one in [Bil97]. Nevertheless, for the canonical height on a projective space, a small sequence of points is strict in our sense if and only if it eventually avoids any fixed translate of a subtorus with essential minimum equal to 0. Such a translate of a subtorus is necessarily a torsion subvariety, see for instance Example 5.16. Hence, a small sequence of points that is strict in the sense of Bilu [Bil97] is also strict in the sense of Definition 5.6. Thus Theorem 5.7 applied to the canonically metrized divisor at infinity on a projective space specializes to [Bil97, Theorem 1.1].

**Remark 5.9.** To the best of our knowledge, even for the canonical metric it is still not known if the equidistribution property for strict sequences holds for the non-Archimedean places of a global field.

The toric Bogomolov conjecture can be stated as follows: let  $X$  be a toric variety and  $D$  an ample toric divisor on  $X$ . Let  $V \subset X_{0, \overline{\mathbb{K}}}$  be a closed subvariety that is not torsion. Then there exists  $\varepsilon > 0$  such that the subset of algebraic points of  $V$  of canonical height bounded above by  $\varepsilon$ , is not dense in  $V$ . Equivalently, if  $V \subset X_{0, \overline{\mathbb{K}}}$  is a closed subvariety such that  $\mu_{\overline{D}}^{\text{ess}}(V) = 0$ , then  $V$  is a torsion subvariety.

This conjecture was proved by Zhang in the number field case [Zha95]. Bilu obtained a proof of Zhang's theorem based on his equidistribution theorem. In what follows, we extend his approach to the general monocritical case over a number field.

Recall that  $X$  denotes a proper toric variety over a number field  $\mathbb{K}$  and  $\overline{D}$  a toric metrized  $\mathbb{R}$ -divisor on  $X$ . For a subvariety  $V \subset X_{\overline{\mathbb{K}}}$ , we set

$$\mu_{\overline{D}}^{\text{abs}}(V) = \inf\{h_{\overline{D}}(x) \mid x \in V(\overline{\mathbb{K}})\}$$

for the absolute minimum of the height function. The fact that  $\overline{D}$  is toric implies

$$\mu_{\overline{D}}^{\text{ess}}(X) = \mu_{\overline{D}}^{\text{abs}}(X_0), \quad (5.2)$$

see [BPS15, Lemma 3.9(2)]. Therefore, for any subvariety  $V \subset X_{0, \overline{\mathbb{K}}}$ ,

$$\mu_{\overline{D}}^{\text{ess}}(V) \geq \mu_{\overline{D}}^{\text{abs}}(V) \geq \mu_{\overline{D}}^{\text{abs}}(X_0) = \mu_{\overline{D}}^{\text{ess}}(X). \quad (5.3)$$

This motivates the following definition.

**Definition 5.10.** A closed subvariety  $V \subset X_{0, \overline{\mathbb{K}}}$  is  $\overline{D}$ -special if

$$\mu_{\overline{D}}^{\text{ess}}(V) = \mu_{\overline{D}}^{\text{ess}}(X).$$

In particular, an algebraic point  $p$  of  $X_0$  is  $\overline{D}$ -special if and only if  $h_{\overline{D}}(p) = \mu_{\overline{D}}^{\text{ess}}(X)$ .

We also propose the following terminology.

**Definition 5.11.** The toric metrized  $\mathbb{R}$ -divisor  $\overline{D}$  satisfies the *Bogomolov property* if every  $\overline{D}$ -special subvariety of  $X_{0, \overline{\mathbb{K}}}$  is a translate of a subtorus.

Note that if  $X$  is of dimension 1, then the Bogomolov property is trivially satisfied for every metrized divisor.

We consider the problem of deciding if a given toric metrized  $\mathbb{R}$ -divisor satisfies the Bogomolov property. The following result corresponds to Theorem 1.4 in the introduction, and shows that the answer is affirmative for monocritical metrics.

**Theorem 5.12.** *Let  $X$  be a proper toric variety over a number field  $\mathbb{K}$  and  $\overline{D}$  a monocritical metrized  $\mathbb{R}$ -divisor on  $X$  with critical point  $\mathbf{u} = (u_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ . Let  $V$  be a  $\overline{D}$ -special subvariety of  $X_{0, \overline{\mathbb{K}}}$ . Then  $V$  is a translate of a subtorus.*

*Furthermore, if  $u_v \in \text{val}_v(\mathbb{T}(\mathbb{K})) \otimes \mathbb{Q}$  for all  $v$ , then  $V$  is the translate of a subtorus by a  $\overline{D}$ -special point.*

Before giving the proof of this theorem, we study special points and, more generally, special translates of subtori in the monocritical case. We first give a criterion for the existence of such points.

**Proposition 5.13.** *Let  $X$  be a proper toric variety over  $\mathbb{K}$  and  $\overline{D}$  a monocritical metrized  $\mathbb{R}$ -divisor on  $X$  with critical point  $\mathbf{u} = (u_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ . Then there exists a  $\overline{D}$ -special point if and only if*

$$u_v \in \text{val}_v(\mathbb{T}(\mathbb{K})) \otimes \mathbb{Q} \quad \text{for all } v \in \mathfrak{M}_{\mathbb{K}}. \quad (5.4)$$

*If this is the case, then every  $\overline{D}$ -special point is of the form  $q^{1/\ell}$  with  $q \in X_0(\mathbb{K})$  and  $\ell \geq 1$ .*

*Proof.* Suppose that there is a  $\overline{D}$ -special point  $p \in X_0(\overline{\mathbb{K}})$ . Choose a finite normal extension  $\mathbb{F} \subset \overline{\mathbb{K}}$  of  $\mathbb{K}$  where  $p$  is defined. Consider the norm of  $p$  relative to this extension, given by

$$N_{\mathbb{K}}^{\mathbb{F}}(p) = \prod_{\tau \in \text{Gal}(\mathbb{F}/\mathbb{K})} \tau(p^{[\mathbb{F}:\mathbb{K}]_i})$$

where  $\text{Gal}(\mathbb{F}/\mathbb{K})$  and  $[\mathbb{F}:\mathbb{K}]_i$  are the Galois group and the inseparable degree of the extension, respectively.

Let  $v \in \mathfrak{M}_{\mathbb{K}}$ . For every  $\tau \in \text{Gal}(\mathbb{F}/\mathbb{K})$ , there is a place  $w \in \mathfrak{M}_{\mathbb{F}}$  over  $v$  such that  $\text{val}_v(\tau(p)) = \text{val}_w(p)$ . By Corollary 4.9 and Proposition 4.17, we have that  $\text{val}_w(p) = u_v$  for any such place. It follows that  $\text{val}_v(\tau(p)) = u_v$  for all  $\tau$ . Using that  $\#\text{Gal}(\mathbb{F}/\mathbb{K}) \cdot [\mathbb{F}:\mathbb{K}]_i = [\mathbb{F}:\mathbb{K}]$ , we deduce that

$$\text{val}_v(N_{\mathbb{K}}^{\mathbb{F}}(p)) = \sum_{\tau} \text{val}_v(\tau(p)^{[\mathbb{F}:\mathbb{K}]_i}) = [\mathbb{F}:\mathbb{K}]u_v.$$

Since  $N_{\mathbb{K}}^{\mathbb{F}}(p) \in \mathbb{T}(\mathbb{K})$ , we get that  $[\mathbb{F}:\mathbb{K}]u_v \in \text{val}_v(\mathbb{T}(\mathbb{K}))$ , proving the implication.

Conversely, assume that the condition (5.4) holds. Let  $S \subset \mathfrak{M}_{\mathbb{K}}$  be a finite set containing the Archimedean places and those places  $v$  where  $u_v \neq 0$ . Set

$$\mathbb{T}(\mathbb{K})_S = \{p \in \mathbb{T}(\mathbb{K}) \mid \text{val}_v(p) = 0 \text{ for all } v \notin S\}$$

and let  $H_{\mathbb{K},S}$  be the subspace of  $\bigoplus_{v \in S} N_{\mathbb{R}}$  defined by the equation  $\sum_{v \in S} n_v z_v = 0$ . Moreover, consider the lattice

$$\Gamma = H_{\mathbb{K},S} \cap \bigoplus_{v \in S} \text{val}_v(\mathbb{T}(\mathbb{K}))$$

and the map  $\text{val}_S: \mathbb{T}(\mathbb{K})_S \rightarrow \Gamma$  given by  $\text{val}_S(p) = (\text{val}_v(p))_{v \in S}$ . By Dirichlet's unit theorem [Wei74, Chapter IV, §4, Corollary to Theorem 9], the image  $\Lambda$  of this map is a sublattice that is commensurable to  $\Gamma$ . Thus  $\Lambda \otimes \mathbb{Q} = \Gamma \otimes \mathbb{Q}$ . Condition (5.4) implies that  $(u_v)_{v \in S} \in \Gamma \otimes \mathbb{Q} = \Lambda \otimes \mathbb{Q}$ . Hence, there is an integer  $\ell \geq 1$  such that

$$(\ell u_v)_{v \in S} \in \Lambda.$$

In other terms, there is  $q \in \mathbb{T}(\mathbb{K})_S$  such that  $\text{val}_v(q) = \ell u_v$  for all  $v \in S$ . By Corollary 4.9, the point  $p = q^{1/\ell} \in \mathbb{T}(\overline{\mathbb{K}})$  is  $\overline{D}$ -special, proving the reverse implication.

To prove the last statement, suppose that the condition (5.4) holds and consider an arbitrary  $\overline{D}$ -special point  $p' \in X_0(\overline{\mathbb{K}})$ . Let  $p$  be the  $\overline{D}$ -special point constructed above and  $\mathbb{F} \subset \overline{\mathbb{K}}$  a finite extension of  $\mathbb{K}$  so that  $p, p' \in \mathbb{T}(\mathbb{F})$ . Then  $\text{val}_w(p'p^{-1}) = 0$  for all  $w \in \mathfrak{M}_{\mathbb{F}}$ . By Kronecker's theorem, the point  $p'p^{-1}$  is torsion. We conclude that some positive power of  $p'$  lies in  $\mathbb{T}(\mathbb{K})$ , as stated.  $\square$

Next we characterize the translates of subtori that are  $\overline{D}$ -special. Let  $U = T_{\overline{\mathbb{K}}} \cdot p$  be the translate of a subtorus  $T \subset \mathbb{T}$  by a point  $p \in X_0(\overline{\mathbb{K}})$ . The subtorus  $T$  corresponds to a saturated sublattice  $Q$  of  $N$ ; we denote by  $\iota: Q \hookrightarrow N$  the corresponding inclusion map. Let  $\mathbb{F} \subset \overline{\mathbb{K}}$  be a finite extension of  $\mathbb{K}$  where  $p$  is defined. For each  $w \in \mathfrak{M}_{\mathbb{F}}$ , we consider the affine subspace of  $N_{\mathbb{R}}$  given by

$$A_{U,w} = \text{val}_w(p) + Q_{\mathbb{R}}.$$

Indeed  $A_{U,w} = \text{val}_w(U_w^{\text{an}})$  and so this affine subspace depends only on  $U$  and not on a particular choice for the translating point  $p$ .

As explained in [BPS14, §3.2], the normalization of the closure of  $U$  in  $X_{\overline{\mathbb{K}}}$  can be given a structure of toric variety. Let  $\Sigma$  be the fan on  $N_{\mathbb{R}}$  corresponding to  $X$  and  $\Sigma_Q$  the fan on  $Q_{\mathbb{R}}$  obtained by restricting  $\Sigma$  to this latter linear space. Then the inclusion  $\iota: Q_{\mathbb{R}} \hookrightarrow N_{\mathbb{R}}$  induces an equivariant map of toric varieties

$$\varphi_{p,\iota}: X_{\Sigma_Q, \overline{\mathbb{K}}} \rightarrow X_{\overline{\mathbb{K}}}$$

extending the inclusion  $U \hookrightarrow \mathbb{T}_{\overline{\mathbb{K}}}$ .

**Proposition 5.14.** *Let  $X$  be a proper toric variety over a number field  $\mathbb{K}$  and  $\overline{D}$  a monocritical metrized  $\mathbb{R}$ -divisor on  $X$  with critical point  $\mathbf{u} = (u_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ . Let  $U = T_{\overline{\mathbb{K}}} \cdot p \subset X_{0, \overline{\mathbb{K}}}$  be the translate of a subtorus  $T \subset \mathbb{T}$  by a point  $p \in X_0(\overline{\mathbb{K}})$  defined over a finite extension  $\mathbb{F} \subset \overline{\mathbb{K}}$  of  $\mathbb{K}$ . For a place  $w$  in  $\mathfrak{M}_{\mathbb{F}}$  denote by  $v(w)$  the place in  $\mathfrak{M}_{\mathbb{K}}$  below  $w$ . Then we have the following properties.*

- (1) *The translate  $U$  is  $\overline{D}$ -special if and only if  $u_{v(w)} \in A_{U,w}$  for all  $w \in \mathfrak{M}_{\mathbb{F}}$ .*
- (2) *If the translate  $U$  is  $\overline{D}$ -special, then the metrized  $\mathbb{R}$ -divisor  $\varphi_{p,\iota}^* \overline{D}$  is monocritical and its critical point is  $(u_{v(w)} - \text{val}_w(p))_{w \in \mathfrak{M}_{\mathbb{F}}}$ .*

*Proof.* By passing to a suitable large finite extension of  $\mathbb{K}$  and applying Proposition 4.17, we can reduce to the case when  $U$  is the translate of a  $\mathbb{K}$ -rational point, that is,  $U = T_{\overline{\mathbb{K}}} \cdot p$  with  $p \in X_0(\mathbb{K})$ . With this assumption,  $\mathbb{F} = \mathbb{K}$  and we set  $v := w = v(w)$ .

Since  $\overline{D}$  is a semipositive toric metrized divisor with  $D$  big, the virtual support function  $\Psi_D$  is concave and its associated polytope has dimension  $n$ . Hence, there is  $m \in M_{\mathbb{R}}$  such that  $\langle m, u \rangle > \Psi_D(u)$  for all  $u \neq 0$ . Moreover, the metric functions  $\psi_{\overline{D},v}$  are concave for all  $v \in \mathfrak{M}_{\mathbb{K}}$ .

Consider the toric metrized  $\mathbb{R}$ -divisor  $\bar{E} := \varphi_{\iota, p}^* \bar{D}$  on the toric variety  $X_{\Sigma_Q}$ . By [BPS14, Proposition 4.3.19], its virtual support function and metric functions are given, for  $z \in Q_{\mathbb{R}}$ , by

$$\Psi_E(z) = \Psi_D(\iota(z)), \quad \psi_{\bar{E}, v}(z) = \psi_{\bar{D}, v}(\text{val}_v(p) + \iota(z)).$$

Therefore  $\Psi_E$  is concave and satisfies  $\langle \iota^\vee m, z \rangle > \Psi_E(z)$  for all  $z \in Q_{\mathbb{R}} \setminus \{0\}$ . Hence, the  $\mathbb{R}$ -divisor  $E$  is big. Moreover, the metric functions  $\psi_{\bar{E}, v}$  are concave and so  $\bar{E}$  is semipositive.

Since  $U$  is identified with a dense open subset of  $X_{\Sigma_Q, \bar{\mathbb{K}}}$ , we have

$$\mu_{\bar{D}}^{\text{ess}}(U) = \mu_{\bar{E}}^{\text{ess}}(X_{\Sigma_Q}).$$

Consider the affine subspace  $\mathbf{A}_U = \bigoplus_v A_{U, v}$  of  $\bigoplus_v N_{\mathbb{R}}$ . By Corollary 4.10,

$$\mu_{\bar{E}}^{\text{ess}}(X_{\Sigma_Q}) = \min_{\mathbf{u}' \in H_{\bar{\mathbb{K}}} \cap \mathbf{A}_U} \sum_v -n_v \psi_{\bar{D}, v}(u'_v), \quad \mu_{\bar{D}}^{\text{ess}}(X) = \min_{\mathbf{u}' \in H_{\bar{\mathbb{K}}}} \sum_v -n_v \psi_{\bar{D}, v}(u'_v).$$

Since  $\bar{D}$  monocritical, the minimum in the right equality is attained only at the point  $\mathbf{u}' = \mathbf{u}$ . We conclude that  $\mu_{\bar{E}}^{\text{ess}}(U) = \mu_{\bar{D}}^{\text{ess}}(X)$  if and only if  $u_v \in A_{U, v}$  for all  $v \in \mathfrak{M}_{\bar{\mathbb{K}}}$ , proving both statements.  $\square$

**Corollary 5.15.** *Let  $X$  be a proper toric variety over a number field  $\mathbb{K}$  and  $\bar{D}$  a monocritical metrized  $\mathbb{R}$ -divisor on  $X$  with critical point  $\mathbf{u} = (u_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ , and suppose that  $u_v \in \text{val}_v(\mathbb{T}(\mathbb{K})) \otimes \mathbb{Q}$  for all  $v \in \mathfrak{M}_{\mathbb{K}}$ . Then a translate of a subtorus of  $X_0$  is  $\bar{D}$ -special if and only if it is the translate of a subtorus by a  $\bar{D}$ -special point.*

*Proof.* Clearly, the translate of a subtorus by a  $\bar{D}$ -special point is  $\bar{D}$ -special. To prove the reverse implication, let  $U$  be a  $\bar{D}$ -special translate of a subtorus and write  $U = T_{\bar{\mathbb{K}}} \cdot p$  as in the statement of Proposition 5.14. By this result, the toric metrized  $\mathbb{R}$ -divisor  $\bar{E} = \varphi_{p, \iota}^* \bar{D}$  is monocritical and, for each  $v \in \mathfrak{M}_{\mathbb{K}}$  and  $w \in \mathfrak{M}_{\mathbb{F}}$  over  $v$ ,

$$u_v \in A_{U, w} \cap \text{val}_v(\mathbb{T}(\mathbb{K})) \otimes \mathbb{Q} \subset A_{U, w} \cap \text{val}_w(\mathbb{T}(\mathbb{F})) \otimes \mathbb{Q}.$$

Since  $p \in X_0(\mathbb{F})$ ,

$$A_{U, w} \cap \text{val}_w(\mathbb{T}(\mathbb{F})) \otimes \mathbb{Q} = \text{val}_w(p) + \text{val}_w(T(\mathbb{F})) \otimes \mathbb{Q}.$$

Hence  $u_v - \text{val}_w(p) \in \text{val}_w(T(\mathbb{F})) \otimes \mathbb{Q}$ . Extending the base field to  $\mathbb{F}$  and restricting to  $X_{\Sigma_Q}$ , Proposition 5.13 implies that this toric variety contains an  $\bar{E}$ -special point. Hence  $U$  contains a  $\bar{D}$ -special point and it is the translate of  $T$  by this point, as stated.  $\square$

**Example 5.16.** Let  $\bar{D}^{\text{can}}$  be a nef and big toric  $\mathbb{R}$ -divisor on the proper toric variety  $X$ , equipped with the canonical metric. By Example 4.15, it is monocritical with critical point  $\mathbf{0} \in H_{\mathbb{K}}$ . Hence,  $p \in X_0(\bar{\mathbb{K}})$  is  $\bar{D}^{\text{can}}$ -special if and only if  $\text{val}_v(p) = 0$  for every  $v \in \mathfrak{M}_{\bar{\mathbb{K}}}$ . By Kronecker's theorem, this is also equivalent to the fact that  $p$  is torsion. Hence, Corollary 5.15 shows that a translate of a subtorus that is  $\bar{D}^{\text{can}}$ -special is necessarily the translate of a subtorus by a torsion point, that is, a torsion subvariety.

*Proof of Theorem 5.12.* Let  $U \subset X_{0, \bar{\mathbb{K}}}$  be the minimal translate of a subtorus containing the subvariety  $V$  and let  $Q$  and  $\Sigma_Q$  be as the ones defined before Proposition 5.14. By (5.2) and (5.3), we have  $\mu_{\bar{D}}^{\text{abs}}(U) = \mu_{\bar{D}}^{\text{ess}}(U)$  and

$$\mu_{\bar{D}}^{\text{ess}}(X) = \mu_{\bar{D}}^{\text{abs}}(X_0) \leq \mu_{\bar{D}}^{\text{abs}}(U) \leq \mu_{\bar{D}}^{\text{abs}}(V) \leq \mu_{\bar{D}}^{\text{ess}}(V) = \mu_{\bar{D}}^{\text{ess}}(X).$$

Therefore,  $U$  is  $\bar{D}$ -special. By Proposition 5.14(2),  $\bar{D}$  pulls back to a monocritical metrized  $\mathbb{R}$ -divisor on  $X_{\Sigma_Q}$ , the normalization of the closure of  $U$  in  $X_{\bar{\mathbb{K}}}$ . Replacing  $X$  by this toric variety, we reduce to the case where  $U = X_{0, \bar{\mathbb{K}}}$ .

Using Proposition 2.5, we choose a sequence  $(p_l)_{l \geq 1}$  of algebraic points of  $V$  that is generic in  $V$  and satisfies

$$\lim_l h_{\overline{D}}(p_l) = \mu_{\overline{D}}^{\text{ess}}(V).$$

Since  $V$  is not contained in any proper translate of a subtorus, this sequence is strict and, since  $V$  is  $\overline{D}$ -special, it is also  $\overline{D}$ -small.

Applying Theorem 5.7 to an Archimedean place  $v \in \mathfrak{M}_{\mathbb{K}}$ , we obtain that the sequence of measures  $(\mu_{p_l, v})_{l \geq 1}$  converges to a measure whose support is the translate  $\mathbb{S}_v \cdot e^{-u_v}$  of the compact subtorus, with  $u_v$  the  $v$ -adic coordinate of the critical point of  $\overline{D}$ .

Since  $V$  is  $\overline{D}$ -special, it is a closed subvariety of  $X_{0, \overline{\mathbb{K}}}$ . Therefore  $V_v^{\text{an}}$  is closed in  $X_{0, v}^{\text{an}}$ . The measures  $(\mu_{p_l, v})_{l \geq 1}$  have support in  $V_v^{\text{an}}$ , and the limit measure has support  $\mathbb{S}_v \cdot e^{-u_v}$ . By the closedness of  $V_v^{\text{an}}$  we deduce the inclusion  $\mathbb{S}_v \cdot e^{-u_v} \subset V_v^{\text{an}}$ . Using that  $\mathbb{S}_v \cdot e^{-u_v}$  is dense in  $X_v^{\text{an}}$  with respect to the Zariski topology, it follows that  $V = X_{0, \overline{\mathbb{K}}}$ , proving the first statement of the theorem.

The last statement of the theorem follows from Corollary 5.15.  $\square$

By Theorem 5.12 and Example 5.16, the canonical toric metrized  $\mathbb{R}$ -divisor  $\overline{D}^{\text{can}}$  satisfies the Bogomolov property, and every  $\overline{D}^{\text{can}}$ -special subvariety is torsion. Hence, Theorem 5.12 extends Zhang's theorem to the general monocritical case. On the other hand, in §6.3 we will give examples of non-monocritical metrized divisors not satisfying the Bogomolov property.

## 6. EXAMPLES

The obtained criteria can be applied in concrete situations to decide if a given semipositive toric metrized  $\mathbb{R}$ -divisor satisfies properties like modulus concentration or equidistribution. In this section, we consider translates of subtori with the canonical height, and toric metrized  $\mathbb{R}$ -divisors equipped with positive smooth metrics at the Archimedean places and canonical metrics at the non-Archimedean ones. We also give a family of counterexamples to the Bogomolov property in the non-monocritical case.

**6.1. Translates of subtori with the canonical height.** Let  $X$  be a proper toric variety of dimension  $n$  over a global field  $\mathbb{K}$  and  $D$  a big and nef toric  $\mathbb{R}$ -divisor on  $X$ . Let  $\Psi_D$  be its virtual support function.

We denote by  $\overline{D}^{\text{can}}$  this  $\mathbb{R}$ -divisor equipped with the canonical metric as in Example 4.1. This toric metrized  $\mathbb{R}$ -divisor satisfies that, for all  $v \in \mathfrak{M}_{\mathbb{K}}$ ,

$$\psi_{\overline{D}^{\text{can}}, v} = \Psi_D \quad \text{and} \quad \vartheta_{\overline{D}^{\text{can}}, v} = 0.$$

Since  $D$  is big,  $\Delta_D$  has dimension  $n$ . Every point  $x$  in the interior of  $\Delta_D$  maximizes the global roof function and  $\partial \vartheta_{\overline{D}^{\text{can}}, v}(x) = \{0\}$ . Therefore, for all  $v \in \mathfrak{M}_{\mathbb{K}}$ ,

$$B_v = \{0\} \quad \text{and} \quad F_v = \{0\}.$$

By Proposition 4.16, the canonical metric is monocritical and so, by Theorem 5.2,  $\overline{D}^{\text{can}}$  satisfies the equidistribution property at every place (Example 5.5).

We next study the toric metrics on  $D$  that are obtained as the inverse image by an equivariant map of a canonical metrized toric divisor on a projective space. For  $r \geq 0$ , let  $\mathbb{P}_{\mathbb{K}}^r$  be the standard projective space over  $\mathbb{K}$  with homogeneous coordinates  $(z_0 : \cdots : z_r)$  and  $H$  the hyperplane at infinity, defined by the equation  $z_0 = 0$ . Denote by  $\overline{H}^{\text{can}}$  this toric divisor equipped with the canonical metric.

Let  $v \in \mathfrak{M}_{\mathbb{K}}$ . If  $v$  is Archimedean, we set  $\lambda_v = 1$  whereas, if  $v$  is non-Archimedean, we set  $\lambda_v$  as the positive generator of the discrete subgroup  $\text{val}_v(\mathbb{K}^\times)$



of  $\mathbb{R}$ . A piecewise affine function is said to be  $\lambda_v$ -rational if all its defining affine functions  $\langle x, u \rangle + b$  satisfy  $x \in M_{\mathbb{Q}}$  and  $b \in \lambda_v \mathbb{Q}$ .

Let  $\psi: N_{\mathbb{R}} \rightarrow \mathbb{R}$  be a concave  $\lambda_v$ -rational piecewise affine function with  $|\psi - \Psi_D|$  bounded. This determines a semipositive metric on  $\mathcal{O}_{X_v^{\text{an}}}(D)$ . As seen in [BPS14, Example 3.7.11], there is an integer  $r > 0$  and a toric morphism  $\iota: X \rightarrow \mathbb{P}_{\mathbb{K}}^r$  such that

$$\psi = \psi_{\iota^* \overline{H}^{\text{can}}, v}.$$

Hence, any such function  $\psi$  can be realized as the  $v$ -adic metric function of the preimage of  $\overline{H}^{\text{can}}$  to  $X$ . This allows us to construct many examples, both monocritical and non-monocritical, of metrized toric divisors.

In the next examples, we fix  $\mathbb{K} = \mathbb{Q}$  and, as before, we denote by  $\overline{H}^{\text{can}}$  the hyperplane at infinity with the canonical metric.

**Example 6.1.** Let  $\iota: \mathbb{G}_{m, \mathbb{Q}} \rightarrow \mathbb{P}_{\mathbb{Q}}^2$  be the map given by

$$\iota(t) = (1 : t/2 : t).$$

Let  $X$  be the normalization of the closure of  $\iota(\mathbb{G}_{m, \mathbb{Q}})$  and  $\overline{D} = \iota^*(\overline{H}^{\text{can}})$ . Then  $X = \mathbb{P}_{\mathbb{Q}}^1$  and  $D$  is the divisor at infinity.

We have  $\Delta_D = [0, 1]$ . As explained in [BPS14, Example 5.1.16], for each  $v \in \mathfrak{M}_{\mathbb{Q}}$  the graph of the local roof function associated to  $\overline{D}$  is given by the upper envelope of the extended polytope

$$\text{conv}((0, 0), (1, \log |1/2|_v), (1, \log |1|_v)) \subset \mathbb{R} \times \mathbb{R}.$$

The graphs of these functions are represented in Figure 1. Thus, for  $x \in [0, 1]$




FIGURE 1. Local roof functions in Example 6.1

we have  $\vartheta_2(x) = x \log(2)$  and  $\vartheta_v(x) = 0$  for  $v \neq 2$ . The global roof function is  $\vartheta(x) = x \log(2)$  and the only point that maximizes it is  $x = 1$ . Moreover,  $\partial \vartheta_2(1) = (-\infty, \log(2)]$  and  $\partial \vartheta_v(1) = (-\infty, 0]$  for  $v \neq 2$ . With Notation 4.2, we have

$$\begin{aligned} B_2 &= [0, \log(2)], & F_2 &= [-\infty, \log(2)], \\ B_v &= [-\log(2), 0], & F_v &= [-\infty, 0] \text{ for } v \neq 2. \end{aligned}$$

By Corollary 4.13, this metrized divisor does not satisfy the modulus concentration property at any place. *A fortiori*, it does not satisfy the equidistribution property at any place.

Indeed, by (4.3) we have  $\mu_{\overline{D}}^{\text{ess}}(X) = \log(2)$ . Let  $(\omega_l)_{l \geq 1}$  be a sequence given by a choice of a primitive  $l$ -th root of the unity,  $a \neq 2$  a positive prime number and  $r$  an integer with  $\log(a) \leq r \log(2)$ . Choose any  $r$ -th root  $a^{1/r}$  of  $a$  and consider the generic sequences of points

$$p_l = (1 : \omega_l) \quad \text{and} \quad q_l = (1 : 2a^{-1/r} \omega_l) \quad \text{for } l \geq 1.$$

For every  $v \in \mathfrak{M}_{\mathbb{Q}}$ ,  $l \geq 1$ ,  $p \in \text{Gal}(p_l)_v$  and  $q \in \text{Gal}(q_l)_v$  we have  $(\text{val}_v)_*(p) = 0$  and

$$(\text{val}_v)_*(q) = \begin{cases} \log(2) & \text{if } v = 2, \\ \frac{-1}{r} \log(a) & \text{if } v = a, \\ -\log(2) + \frac{1}{r} \log(a) & \text{if } v = \infty, \\ 0 & \text{if } v \neq 2, a, \infty. \end{cases}$$

Either by computing the local roof functions of  $\overline{D}$  or the Weil height of the image of these points under the inclusion  $\iota$ , we deduce that

$$h_{\overline{D}}(p_l) = \log(2) \quad \text{and} \quad h_{\overline{D}}(q_l) = \log(2).$$

Therefore both sequences are  $\overline{D}$ -small. For any place  $v$ , the sequence  $\mu_{p_l, v}$  converges to  $\lambda_{\mathbb{S}_v, 0}$ . In contrast, if we denote  $u_v = (\text{val}_v)_*(q)$  for any  $q \in \text{Gal}(q_l)_v$ , then  $\mu_{q_l, v}$  converges to  $\lambda_{\mathbb{S}_v, u_v}$ . This shows that neither the modulus concentration nor the equidistribution properties hold for the places  $2, a, \infty$ . Varying  $a$ , we deduce that these properties do not hold at any place of  $\mathbb{Q}$ .

The metric of  $\overline{D}$  at the Archimedean place is the canonical one. The metrics at the non-Archimedean places can be interpreted in terms of integral models. Let  $\mathcal{X}$  be the blow up of  $\mathbb{P}_{\mathbb{Z}}^1$  at the point  $(1 : 0)$  over the prime 2. The fibre of the structural map  $\mathcal{X} \rightarrow \text{Spec}(\mathbb{Z})$  over the point 2 has two components: the exceptional divisor of the blow up, which we denote by  $E$ , and the strict transform of the fibre of  $\mathbb{P}_{\mathbb{Z}}^1$ , which we denote by  $Y$ . Consider the divisor

$$\mathcal{D} = \overline{\infty} + Y,$$

where  $\overline{\infty}$  denotes the closure in  $\mathcal{X}$  of the point  $(0 : 1) \in \mathbb{P}^1(\mathbb{Q})$ . The pair  $(\mathcal{X}, \mathcal{D})$  is a model of  $(X, D)$ . For each non-Archimedean place  $v$ , this model induces an algebraic metric on  $D$  that agrees with the  $v$ -adic metric of  $\overline{D}$ .

**Example 6.2.** Consider now the map  $\iota: \mathbb{G}_{m, \mathbb{Q}} \rightarrow \mathbb{P}_{\mathbb{Q}}^2$  given by

$$\iota(t) = (t^{-1} : 1/2 : t).$$

Let  $X$  be the normalization of the closure of  $\iota(\mathbb{G}_{m, \mathbb{Q}})$  and  $\overline{D} = \iota^*(\overline{H}^{\text{can}})$ . In this case,  $X = \mathbb{P}_{\mathbb{Q}}^1$  and  $D$  is the divisor at infinity plus the divisor at zero.

We have  $\Delta_D = [-1, 1]$ . As before, we compute the local roof functions using [BPS14, Example 5.1.16]. Their graphs are represented in Figure 2. For  $x \in [0, 2]$ ,




FIGURE 2. Local roof functions in Example 6.2

we have  $\vartheta_2(x) = (1 - |x|) \log(2)$  and  $\vartheta_v(x) = 0$  for  $v \neq 2$ . Thus, the global roof function is  $\vartheta(x) = (1 - |x|) \log(2)$ . Its maximum is attained only at the point  $x = 0$ . In this case,  $\partial\vartheta_2(0) = [-\log(2), \log(2)]$  and  $\partial\vartheta_v(0) = \{0\}$  for  $v \neq 2$ . We deduce that

$$B_2 = \{0\}, F_2 = [-\log(2), \log(2)] \quad \text{and} \quad B_v = \{0\}, F_v = \{0\} \text{ for } v \neq 2. \quad (6.1)$$

By Corollary 4.13,  $\overline{D}$  satisfies modulus concentration for all places except the place 2. This toric metrized divisor is not monocritical, and so we cannot apply Theorem 5.2 in this case. Indeed, later we will see that  $\overline{D}$  does not satisfy the equidistribution property at any place of  $\mathbb{Q}$  (Example 7.6).

As in the previous example, the metric of  $\overline{D}$  at the Archimedean place is the canonical one, and those at the non-Archimedean places can be interpreted in terms of integral models. Let  $\mathcal{X}$  be the blow up of  $\mathbb{P}_{\mathbb{Z}}^1$  at the points  $(1 : 0)$  and  $(0 : 1)$  over the prime 2. The fibre of the structural map  $\mathcal{X} \rightarrow \text{Spec}(\mathbb{Z})$  over the point 2 has three components. Consider the divisor

$$\mathcal{D} = \overline{\infty} + \overline{0},$$

where  $\overline{\infty}$  denotes the closure in  $\mathcal{X}$  of the point  $(0 : 1) \in \mathbb{P}^1(\mathbb{Q})$  and  $\overline{0}$  the closure of the point  $(1 : 0)$ . The pair  $(\mathcal{X}, \mathcal{D})$  is a model of  $(X, D)$ . For each non-Archimedean place  $v$ , this model induces an algebraic metric on  $D$  that agrees with the  $v$ -adic metric of  $\overline{D}$ .

**Example 6.3.** This time we consider the map  $\iota : \mathbb{G}_{m, \mathbb{Q}} \rightarrow \mathbb{P}_{\mathbb{Q}}^3$  given by

$$\iota(t) = (1 : t/2 : t^2/2 : t^3).$$

Let  $X$  be the normalization of the closure of  $\iota(\mathbb{G}_{m, \mathbb{Q}})$  and  $\overline{D} = \iota^*(\overline{H}^{\text{can}})$ . In this case,  $X = \mathbb{P}_{\mathbb{Q}}^1$  and  $D$  is three times the divisor at infinity.

We have  $\Delta_D = [0, 3]$  and the local roof functions are represented in Figure 3. They are given by  $\vartheta_2(x) = \log(2) \min(x, 1, 3 - x)$  and  $\vartheta_v(x) = 0$  for  $v \neq 2$ . The




FIGURE 3. Local roof functions in Example 6.2

global roof function is thus  $\vartheta(x) = \log(2) \min(x, 1, 3 - x)$ , which is maximized at any point of the interval  $[1, 2]$ . Choosing the maximizing point  $x = 3/2$ , we have  $\partial\vartheta_v(3/2) = \{0\}$  for all  $v$ .

Thus  $\overline{D}$  is monocritical, by Proposition 4.16. By Corollary 4.13 and Theorem 5.2, it satisfies both the modulus concentration and the equidistribution properties for any place.

**6.2. Positive Archimedean metrics.** The following result covers many of the examples considered in [BPS14, BMPS16, BPS15]: twisted Fubini-Study metrics on projective spaces, metrics from polytopes, Fubini-Study metrics on toric bundles,  $\ell^p$ -metrics on toric varieties, and Fubini-Study metrics on weighted projective spaces. All of them consist of toric varieties over  $\mathbb{Q}$  with a toric divisor equipped with a positive smooth metric at the Archimedean place and the canonical metric at the non-Archimedean ones.

**Theorem 6.4.** *Let  $X$  be a proper toric variety over a number field  $\mathbb{K}$  and  $\overline{D} = (D, (\|\cdot\|_v)_{v \in \mathfrak{M}_{\mathbb{K}}})$  a semipositive toric metrized  $\mathbb{R}$ -divisor with  $D$  big. We assume that, when  $v$  is Archimedean,  $\|\cdot\|_v$  is a positive smooth metric on the principal open subset  $X_{0,v}^{\text{an}}$  whereas, when  $v$  is non-Archimedean, it is the  $v$ -adic canonical metric of  $D$ . Then  $\overline{D}$  is monocritical. In particular, it satisfies the equidistribution property for every place of  $\mathbb{K}$ .*

*When  $\mathbb{K} = \mathbb{Q}$ , the  $v$ -adic limit measure is  $\lambda_{\mathbb{S}_v,0}$  for every  $v \in \mathfrak{M}_{\mathbb{Q}}$ .*

*Proof.* Since the metric is smooth and positive on  $X_{0,v}^{\text{an}}$  for  $v$  Archimedean, the proof of [BPS14, Proposition 4.4.1] implies that the metric function  $\psi_{\overline{D},v}$  is smooth and strictly concave, in the sense that its Hessian is negative definite. Therefore  $\psi_{\overline{D},v}$  is of Legendre type in the sense of [BPS14, Definition 2.4.1] and, by [BPS14, Theorem 2.4.2(2)], the local roof function  $\vartheta_{\overline{D},v}$  is of Legendre type. In particular,  $\vartheta_{\overline{D},v}$  is smooth and strictly concave on the interior of  $\Delta_D$  and the sup-differential at any point of the border of the polytope is empty.

For the non-Archimedean places, the metrics are canonical and so their local roof functions are zero. Hence

$$\vartheta_{\overline{D}} = \sum_{v|\infty} n_v \vartheta_{\overline{D},v},$$

this function is smooth and strictly concave on the interior of  $\Delta_D$ , and its sup-differential at any point of the border of  $\Delta_D$  is empty. This implies that there is a unique maximizing point  $x_{\max} \in \Delta_D$ , which lies in the interior of the polytope, and that  $\partial \vartheta_{\overline{D}}(x_{\max}) = \{0\}$ . Thus, the first assertion then follows from Proposition 4.16.

When  $\mathbb{K} = \mathbb{Q}$  there is only one Archimedean place. Therefore all the  $v$ -adic metrics are the canonical metric except one. This implies easily that the critical point in this case is  $\mathbf{u} = (0)_v$  and the last statement follows from Theorem 5.2.  $\square$

**Example 6.5.** Let  $X = \mathbb{P}_{\mathbb{Q}}^1$  and  $\overline{D}$  the divisor at infinity equipped with the Fubini-Study metric at the Archimedean place and the canonical metric at the non-Archimedean ones. By Theorem 6.4, this toric metrized divisor satisfies the equidistribution property at every place. Moreover, the limit measure of the Galois orbits of any generic  $\overline{D}$ -small sequence is  $\lambda_{\mathbb{S}_v,0}$ .

Recall that the canonical metric at the non-Archimedean places corresponds to the canonical model of  $(\mathbb{P}_{\mathbb{Q}}^1, \infty)$  given by  $(\mathbb{P}_{\mathbb{Z}}^1, \overline{\infty})$ , where  $\overline{\infty}$  is the closure of the point  $(0 : 1) \in \mathbb{P}^1(\mathbb{Q})$ . If we change the integral model, different phenomena may occur. For instance, consider the integral model of Example 6.1, whose global roof function is given by

$$\vartheta_{\overline{D},\infty}(x) = -\frac{1}{2}(x \log x + (1-x) \log(1-x)) + x \log(2),$$

see [BPS14, Example 6.2.3]. The unique maximum of this function is attained at a point in the interior of  $\Delta_D = [0, 1]$ . Since  $\vartheta_{\overline{D},\infty}$  is differentiable on  $(0, 1)$ , we deduce that the sup-differential is reduced to one point. By Proposition 4.16, this new toric metrized divisor is also monocritical.

In contrast, if we consider the divisor  $D' = 0 + \infty$  with the Fubini-Study metric at the Archimedean place and the metrics induced by the integral model of Example 6.2, then the maximum of the global roof function is attained at the point zero and the sup-differential at this point is  $[-\log(2), \log(2)]$ . Since zero is not a vertex of this set, by Proposition 4.16 this divisor is not monocritical. Hence it does not satisfy the equidistribution property at the Archimedean place.

**6.3. Counterexamples to the Bogomolov property.** In this section, we give examples of toric metrized divisors not satisfying the Bogomolov property. For

simplicity, we restrict to the case  $\mathbb{K} = \mathbb{Q}$ . As in §6.1, we denote by  $\overline{H}^{\text{can}}$  the canonical metrized divisor at infinity on a projective space.

**Example 6.6.** Consider the map  $\iota: \mathbb{G}_{m,\mathbb{Q}} \times \mathbb{G}_{m,\mathbb{Q}} \rightarrow \mathbb{P}_{\mathbb{Q}}^3$  given by

$$\iota(t_1, t_2) = (1 : 2 : t_1 : t_2).$$

As in the examples in the previous section, we denote by  $X$  the normalization of the closure of the image of  $\iota$  and  $\overline{D} = \iota^*(\overline{H}^{\text{can}})$ . In this case,  $X = \mathbb{P}_{\mathbb{Q}}^2$  and  $D$  is the divisor at infinity.

We have that  $\Delta_D$  is the standard simplex of  $N_{\mathbb{R}} = \mathbb{R}^2$  and  $\Psi_D: \mathbb{R}^2 \rightarrow \mathbb{R}$  is the function given by

$$\Psi_D(u_1, u_2) = \min(0, u_1, u_2).$$

By [BPS14, Example 4.3.21], the local metric functions are given, for  $(u_1, u_2) \in \mathbb{R}^2$ , by

$$\psi_{\overline{D},v}(u_1, u_2) = \begin{cases} \Psi_D(u_1 + \log(2), u_2 + \log(2)) - \log(2) & \text{if } v = \infty, \\ \Psi_D(u_1, u_2) & \text{if } v \neq \infty. \end{cases}$$

By [BPS14, Example 5.1.16], the local roof functions are given, for  $(x_1, x_2) \in \Delta_D$ , by

$$\vartheta_{\overline{D},v}(x_1, x_2) = \begin{cases} (1 - x_1 - x_2) \log(2) & \text{if } v = \infty, \\ 0 & \text{if } v \neq \infty. \end{cases}$$

Hence the global roof function agrees with  $\vartheta_{\overline{D},\infty}$ . Its only maximizing point is  $x_{\max} = (0, 0)$ , and one computes easily that  $\partial\vartheta_{\overline{D},\infty}(0, 0) = (-\log(2), -\log(2)) + \mathbb{R}_{\geq 0}^2$  and  $\partial\vartheta_{\overline{D},v}(0, 0) = \mathbb{R}_{\geq 0}^2$  for  $v \neq \infty$ . Thus

$$B_{\infty} = [-\log(2), 0]^2, \quad F_{\infty} = (-\log(2), -\log(2)) + \mathbb{R}_{\geq 0}^2, \\ B_v = [0, \log(2)]^2, \quad F_v = \mathbb{R}_{\geq 0}^2 \quad \text{for } v \neq \infty.$$

We also have  $\mu_{\overline{D}}^{\text{ess}}(X) = \vartheta(0, 0) = \log(2)$ .

Let  $(z_0 : z_1 : z_2)$  be homogeneous coordinates of  $X$  and consider the curve  $C$  of equation  $z_0 + z_1 + z_2 = 0$ . In what follows, we will see that this curve is a  $\overline{D}$ -special subvariety. Since  $C$  is not a translate of a subtorus, this will show that  $\overline{D}$  does not satisfy the Bogomolov property.

For  $l \geq 1$  choose a primitive  $l$ -th root of the unity  $\omega_l$ . Let  $z_{1,l}$  be a solution of the equation  $z^2 + z + \omega_l = 0$  and put  $z_{2,l} = \omega_l/z_{1,l}$  for the other solution. Then

$$z_{1,l} + z_{2,l} + 1 = 0 \quad \text{and} \quad z_{1,l}z_{2,l} = \omega_l. \quad (6.2)$$

In particular,  $p_l = (1 : z_{1,l} : z_{2,l})$  is an algebraic point of  $C$ .

Let  $v \in \mathfrak{M}_{\mathbb{Q}}$  and  $q = (1 : q_1 : q_2) \in \text{Gal}(p_l)_v$ . If  $v \neq \infty$ , then the conditions (6.2) imply that

$$\text{val}_v(q) = (0, 0) \in B_v. \quad (6.3)$$

If  $v = \infty$ , then these same conditions (6.2) give  $\max(|q_1|_{\infty}, |q_2|_{\infty}) \leq \frac{1+\sqrt{5}}{2}$ . Thus

$$\text{val}_{\infty}(q) \in \left( -\log\left(\frac{1+\sqrt{5}}{2}\right), -\log\left(\frac{1+\sqrt{5}}{2}\right) \right) + \mathbb{R}_{\geq 0}^2 \subset F_{\infty}. \quad (6.4)$$

Moreover, by the product formula and (6.3), we have

$$E[\nu_{p_l, \infty}] = \frac{1}{\#\text{Gal}(p_l)_{\infty}} \sum_{q \in \text{Gal}(p_l)_{\infty}} \text{val}_{\infty}(q) = (0, 0) \in B_{\infty}. \quad (6.5)$$

By Corollary 4.9, the conditions (6.3), (6.4) and (6.5) imply that  $h_{\overline{D}}(p_l) = \mu_{\overline{D}}^{\text{ess}}(X)$ . Since the sequence  $(p_l)_{l \geq 1}$  is generic in  $C$ , we deduce  $\mu_{\overline{D}}^{\text{ess}}(C) = \mu_{\overline{D}}^{\text{ess}}(X)$  and so  $C$  is a  $\overline{D}$ -special subvariety.

We generalize this example to a family of metrics on toric varieties of dimension greater than or equal to 2.

**Proposition 6.7.** *Let  $X$  be a proper toric variety over  $\mathbb{Q}$  of dimension  $n \geq 2$  and  $D$  a big and nef  $\mathbb{R}$ -divisor on  $X$ . Let  $u_0 \in N_{\mathbb{R}}$  and consider the metrized divisor  $\overline{D}^{u_0}$  over  $D$  defined by*

$$\psi_{\overline{D}^{u_0}, v}(u) = \begin{cases} \Psi_D(u - u_0) & \text{if } v = \infty, \\ \Psi_D(u) & \text{if } v \neq \infty. \end{cases}$$

Then  $\overline{D}^{u_0}$  satisfies the Bogomolov property if and only if  $u_0 = 0$ .

*Proof.* When  $u_0 = 0$  we have  $\overline{D}^{u_0} = \overline{D}^{\text{can}}$ . By Theorem 5.12 and Example 5.16, this toric metrized divisor satisfies the Bogomolov property.

Suppose  $u_0 \neq 0$ . The local roof functions of  $\overline{D}^{u_0}$  are given, for  $x \in \Delta_D$ , by

$$\vartheta_{\overline{D}^{u_0}, v}(x) = \begin{cases} \langle x, u_0 \rangle & \text{if } v = \infty, \\ 0 & \text{if } v \neq \infty. \end{cases}$$

In particular, the global roof function  $\vartheta_{\overline{D}}$  coincides with  $\vartheta_{\overline{D}^{u_0}, \infty}$ . The maximum of  $\vartheta_{\overline{D}}$  is attained on a face of  $\Delta_D$ . Fix  $x_0$  in the relative interior of this face. If we denote by  $\vartheta_0$  the constant function equal to 0 defined on  $\Delta_D$ , then  $\sigma_0 = \partial\vartheta_0(x_0)$  is a cone in  $N_{\mathbb{R}}$  containing  $-u_0$  in its relative interior. Moreover,

$$\partial\vartheta_{\overline{D}^{u_0}, \infty}(x_0) = u_0 + \sigma_0 \quad \text{and} \quad \partial\vartheta_{\overline{D}^{u_0}, v}(x_0) = \sigma_0 \text{ for } v \neq \infty.$$

It follows that  $0 \in B_v$  for every  $v$ , that  $F_{\infty} = u_0 + \sigma_0$  and that  $F_v = \sigma_0$  for  $v \neq \infty$ .

As in Example 6.6, to prove that  $\overline{D}^{u_0}$  does not satisfy the Bogomolov property, it is enough to exhibit a curve  $C$  in  $X$  that is  $\overline{D}$ -special but not a translate of a subtorus.

We identify  $N_{\mathbb{R}} \simeq \mathbb{R}^n$ . Since  $X$  is proper and  $\sigma_0$  is a cone of the fan of  $X$ , there is a primitive vector  $n_0 \in N$  in  $\sigma_0$ . It follows that there is  $\varepsilon_0 > 0$  such that

$$\ell_0 := \{xn_0 \mid -\varepsilon_0 \leq x \leq \varepsilon_0\} \subset u_0 + \sigma_0.$$

Choose a primitive vector  $a_0 \in N$  such that  $a_0$  and  $n_0$  generate a saturated sublattice  $V$  of  $N$ . Put  $b_0 = n_0 + a_0$ . Then  $a_0$  and  $b_0$  form an integral basis of  $V$ . Fix an integer  $k_0 \geq \varepsilon_0^{-1}$  and consider the linear map  $L: V_{\mathbb{R}} \rightarrow \mathbb{R}^2$  defined by

$$L(sa_0 + tb_0) = k_0 \cdot (s, t).$$

Let  $S$  be the toric surface in  $X_0$  associated to the saturated sublattice  $V$ . The linear map  $L$  induces a toric morphism  $\iota: S \rightarrow \mathbb{G}_{m, \mathbb{Q}}^2$ . Let  $C$  be the curve in  $\mathbb{G}_{m, \mathbb{Q}}^2$  of equation  $x + y + 1 = 0$  and denote by  $C_0$  the closure in  $X$  of the curve  $\iota^{-1}(C)$ .

As in Example 6.6, for  $l \geq 1$  choose a primitive  $l$ -th root of unity  $\omega_l$ . Let  $z_{1,l}$  be a solution of the equation  $z^2 + z + \omega_l = 0$  and put  $z_{2,l} = \omega_l/z_{1,l}$ . Hence

$$z_{1,l} + z_{2,l} + 1 = 0 \quad \text{and} \quad z_{1,l}z_{2,l} = \omega_l.$$

In particular,  $(z_{1,l}, z_{2,l}) \in C(\overline{\mathbb{Q}})$ . Choose a point  $p_l \in C_0(\overline{\mathbb{Q}})$  such that  $\iota(p_l) = (z_{1,l}, z_{2,l})$ . The sequence of points  $(p_l)_{l \geq 0}$  is generic in  $C_0$ .

For every place  $v$  there is a commutative diagram

$$\begin{array}{ccccc} (\mathbb{G}_m^2)_v^{\text{an}} & \xleftarrow{\iota} & S_v^{\text{an}} & \hookrightarrow & X_{0,v}^{\text{an}} \\ \downarrow \text{val}_v & & \downarrow \text{val}_v & & \downarrow \text{val}_v \\ \mathbb{R}^2 & \xleftarrow{L} & V_{\mathbb{R}} & \hookrightarrow & N_{\mathbb{R}} \end{array}$$

Since  $n_0 = b_0 - a_0$ , we have

$$\ell := L(\ell_0) = \{(x, -x) \mid |x| \leq \varepsilon_0 k_0\}.$$

Arguing as in Example 6.6, for every non-Archimedean place  $v$  and every point  $q \in \text{Gal}(p_l)_v$ , we have

$$\text{val}_v(\iota(q)) = 0.$$

Since  $L$  is injective,  $\text{val}_v(q) = 0$  and therefore  $\nu_{p_l, v} = \delta_0$ . In particular,

$$\text{supp}(\nu_{p_l, v}) = \{0\} \subset F_v \text{ and } E[\nu_{p_l, v}] = 0 \in B_v.$$

When  $v = \infty$ , the product formula implies that

$$E[\nu_{p_l, \infty}] = \frac{1}{\#\text{Gal}(p_l)_\infty} \sum_{q \in \text{Gal}(p_l)_\infty} \text{val}_\infty(q) = 0 \in B_\infty.$$

On the other hand, note that for every  $q$  in  $\text{Gal}(p_l)_\infty$ , the point  $\iota(q) = (q_1, q_2)$  satisfies

$$q_1 + q_2 + 1 = 0 \quad \text{and} \quad q_1 q_2 = \omega_l.$$

We thus have  $|q_1|_\infty |q_2|_\infty = 1$ ,

$$\frac{\sqrt{5} - 1}{2} \leq \min\{|q_1|_\infty, |q_2|_\infty\} \leq \max\{|q_1|_\infty, |q_2|_\infty\} \leq \frac{1 + \sqrt{5}}{2},$$

and therefore

$$|\max(-\log |q_1|_\infty, -\log |q_2|_\infty)| \leq \log \left( \frac{1 + \sqrt{5}}{2} \right) \leq 1 \leq \varepsilon_0 k_0.$$

This implies that

$$\text{val}_\infty(\iota(q)) \in \ell, \quad \text{val}_\infty(q) \in \ell_0 \subset u_0 + \sigma_0 = F_\infty \quad \text{and} \quad \text{supp}(\nu_{p_l, \infty}) \subset F_\infty.$$

By Lemma 4.8, we have  $h_{\overline{D}}(p_l) = \mu_{\overline{D}}^{\text{ess}}(X)$ . Being the sequence  $(p_l)_{l \geq 1}$  generic in  $C_0$ , we deduce that  $C_0$  is  $\overline{D}$ -special. Since  $C_0$  is not a translate of a subtorus, we conclude that  $\overline{D}$  does not satisfy the Bogomolov property, as stated.  $\square$

## 7. POTENTIAL THEORY ON THE PROJECTIVE LINE AND SMALL POINTS

In this section, we apply potential theory on the projective line over a number field, and in particular Rumely's Fekete-Szegő theorem, to produce interesting sequences of small points in the non-monocritical case.

In the absence of modulus concentration, this allows to produce a wealth of non-toric measures that are limit measures of Galois orbits of generic sequences of points of small height. These techniques also allow to show that the absence of modulus concentration at a place can affect the equidistribution property at another place.

**7.1. Limit measures in the absence of modulus concentration.** We recall the basic objects of potential theory on the projective line. For most of the details and precise definitions, we refer the reader to [Tsu75] and [BR10] for the Archimedean and non-Archimedean cases, respectively.

Let  $\mathbb{K}$  be a number field and fix a place  $v \in \mathfrak{M}_{\mathbb{K}}$ . For a subset  $E \subset \mathbb{C}_v$ , we denote by  $\overline{E}$  its closure in  $\mathbb{A}_v^{1, \text{an}}$ . Moreover, for  $r > 0$ , put

$$\mathcal{B}_v(E, r) = \left\{ z \in \mathbb{C}_v \mid \inf_{a \in E} |z - a|_v \leq r \right\}.$$

In particular, for  $a \in \mathbb{C}_v$  the set  $\mathcal{B}_v(a, r)$  is the closed ball with center  $a$  and radius  $r$ . Set  $O_v = \mathcal{B}_v(0, 1)$ , and recall that  $\mathbb{S}_v = \{z \in \mathbb{C}_v \mid |z|_v = 1\}$ .

Note that if  $E$  is a bounded subset of  $\mathbb{C}_v$ , then  $\overline{E}$  is compact. Since  $\mathbb{A}_v^{1, \text{an}}$  is metrizable, it follows that the set of Borel probability measures on  $\overline{E}$  endowed with the weak-\* topology is compact, metrizable, and therefore sequentially compact.

Denoting by  $\mathbb{A}_v^{1, \text{an}} \times \mathbb{A}_v^{1, \text{an}}$  the product of  $\mathbb{A}_v^{1, \text{an}}$  with itself in the category of topological spaces, let

$$\delta_v: \mathbb{A}_v^{1, \text{an}} \times \mathbb{A}_v^{1, \text{an}} \rightarrow \mathbb{R}$$

be the function defined by  $\delta_v(z, z') = |z - z'|_v$  for  $v$  Archimedean, and the unique upper semicontinuous extension of the function on  $\mathbb{C}_v \times \mathbb{C}_v$  defined by  $(z, z') \mapsto |z - z'|_v$  for  $v$  non-Archimedean, see [BR10, Proposition 4.1].

Given a Borel probability measure  $\mu$  on  $\mathbb{A}_v^{1,\text{an}}$ , the *energy integral* (with respect to the point at infinity) of  $\mu$  is defined as

$$I_v(\mu) = \int_{\mathbb{A}_v^{1,\text{an}}} \int_{\mathbb{A}_v^{1,\text{an}}} -\log(\delta_v(z, z')) \, d\mu(z) \, d\mu(z'). \quad (7.1)$$

Let  $K \subset \mathbb{A}_v^{1,\text{an}}$  be a measurable subset. The *v-adic Robin constant* and *capacity* (with respect to the point at infinity) of  $K$  are respectively defined as

$$V_v(K) = \inf\{I_v(\mu) \mid \text{supp}(\mu) \subset K\} \quad \text{and} \quad \text{cap}_v(K) = e^{-V_v(K)}. \quad (7.2)$$

If  $K$  is compact and  $\text{cap}_v(K) > 0$ , then there exists a unique probability measure, denoted by  $\rho_K$ , supported on  $K$  and realizing the infimum in (7.2), see [Tsu75, §III.2 and Theorem III.32] for the Archimedean case and [BR10, Propositions 6.6 and 7.21] for the non-Archimedean one. Hence

$$I_v(\rho_K) = V_v(K).$$

This measure is called the *equilibrium measure* of  $K$ . It does not charge singletons, so we can also consider it as a measure on  $\mathbb{C}_v^\times$ . For  $K = \overline{O}_v$ , it agrees with  $\lambda_{\mathbb{S}_v,0}$ , the Haar probability measure on the unit circle when  $v$  is Archimedean, and the Dirac measure at the Gauss point of  $\mathbb{A}_v^{1,\text{an}}$  when  $v$  is non-Archimedean. We also have

$$\text{cap}_v(O_v) = 1, \quad (7.3)$$

see for example [Rum02, §3].

In the non-Archimedean case,  $\mathbb{C}_v$  is a proper subset of  $\mathbb{A}_v^{1,\text{an}}$ . In general, for a Borel subset  $E$  of  $\mathbb{C}_v$ , we have

$$\text{cap}_v(E) \leq \text{cap}_v(\overline{E}),$$

but this inequality might be strict even if  $E$  is closed and bounded. Equality holds if, for example, there are  $r > 0$  and a polynomial  $P$  with coefficients in  $\mathbb{C}_v$ , such that  $E = \{z \in \mathbb{C}_v \mid |P(z)| \leq r\}$ , see [BR10, Corollary 6.26] and [Rum02, §3.2].

**Definition 7.1.** An *adelic set* is a collection  $\mathbf{E} = (E_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$  such that  $E_v$  is a subset of  $\mathbb{C}_v$  invariant under the action of the absolute  $v$ -adic Galois group  $\text{Gal}(\overline{\mathbb{K}}_v/\mathbb{K}_v)$  for all  $v$ , and such that  $E_v = O_v$  for all but a finite number of  $v$ . We say that  $\mathbf{E}$  is *bounded* (respectively *closed*, *open*) if  $E_v$  is bounded (respectively closed, open) for all  $v$ .

Given an adelic set  $\mathbf{E} = (E_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$ , its (*global*) *capacity* is defined as

$$\text{cap}(\mathbf{E}) = \prod_{v \in \mathfrak{M}_{\mathbb{K}}} \text{cap}_v(E_v)^{n_v}.$$

By (7.3), this product actually runs over a finite set and so the global capacity is well-defined.

The following result shows that, in the non-monocritical case, there is a wealth of limit measures of Galois orbits of generic sequences of points of small height that are not invariant under the action of the compact torus.

**Theorem 7.2.** *Let  $X = \mathbb{P}_{\mathbb{K}}^1$  and  $\overline{D}$  the divisor at infinity equipped with a semipositive toric metric. Let  $B_v, F_v$  be the associated subsets of  $N_{\mathbb{R}} = \mathbb{R}$  as in (4.4). Let  $\mathbf{E} = (E_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$  be a closed bounded adelic set such that  $\text{cap}(\mathbf{E}) = 1$ , and such that for every non-Archimedean place  $v$  we have  $\text{cap}(\overline{E}_v) = \text{cap}(E_v)$ . Assume that the following conditions hold:*

- (1)  $\text{supp}((\text{val}_v)_* \rho_{E_v}) \subset F_v$  for all  $v \in \mathfrak{M}_{\mathbb{K}}$ ;



- (2)  $E[(\text{val}_v)_* \rho_{E_v}] \in B_v$  for all  $v \in \mathfrak{M}_{\mathbb{K}}$ ;  
(3)  $\sum_{v \in \mathfrak{M}_{\mathbb{K}}} n_v E[(\text{val}_v)_* \rho_{E_v}] = 0$ .

Then there is a generic  $\bar{D}$ -small sequence  $(p_l)_{l \geq 1}$  of algebraic points of  $X_0 = \mathbb{G}_{m, \mathbb{K}}$  such that, for every  $v \in \mathfrak{M}_{\mathbb{K}}$ , the sequence of probability measures  $(\mu_{p_l, v})_{l \geq 1}$  converges to  $\rho_{\bar{E}_v}$ .

The proof of this theorem will be given after two preliminary propositions. The next statement is a direct consequence of Rumely's version of the Fekete-Szegő theorem in [Rum02, Theorem 2.1].

**Proposition 7.3.** *Let  $\mathbf{E} = (E_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$  be a closed bounded adelic set such that  $\text{cap}(\mathbf{E}) \geq 1$ . There exists a sequence  $(p_l)_{l \geq 1}$  of pairwise distinct points of  $\bar{\mathbb{K}}^\times$  satisfying*

$$\text{Gal}(p_l)_v \subset \mathcal{B}_v\left(E_v, \frac{1}{l}\right)$$

for all  $l \geq 1$  and  $v \in \mathfrak{M}_{\mathbb{K}}$ . In particular,  $\text{Gal}(p_l)_v \subset E_v$  for every non-Archimedean place  $v$  such that  $E_v = O_v$ .

*Proof.* For  $l \geq 1$ , consider the bounded adelic neighbourhood  $\mathbf{U}_l = (U_{l, v})_{v \in \mathfrak{M}_{\mathbb{K}}}$  of  $\mathbf{E}$  given by

$$U_{l, v} = \mathcal{B}_v\left(E_v, \frac{1}{l}\right).$$

By [Rum02, Theorem 2.1] with  $S = \emptyset$ , there is an infinite number of points  $p \in \bar{\mathbb{K}}^\times$  such that  $\text{Gal}(p)_v \subset U_{l, v}$  for all  $v$ . Inductively, for each  $l \geq 1$  we choose  $p_l$  as one of these points that is different from  $p_{l'}$  for  $l' \leq l - 1$ .  $\square$

In the notation of Proposition 7.3, when the adelic set  $\mathbf{E}$  has capacity 1, the sequence of  $v$ -adic Galois orbits of the points  $p_l$  equidistribute according to the equilibrium measure of the closure  $\bar{E}_v$ .

**Proposition 7.4.** *Let  $\mathbf{E} = (E_v)_{v \in \mathfrak{M}_{\mathbb{K}}}$  be a closed bounded adelic set such that  $\text{cap}(\mathbf{E}) = 1$  and such that for every non-Archimedean place  $v$  we have  $\text{cap}(\bar{E}_v) = \text{cap}(E_v)$ . Let  $(p_l)_{l \geq 1}$  be a sequence of pairwise distinct points of  $\bar{\mathbb{K}}^\times$  with  $\text{Gal}(p_l)_v \subset \mathcal{B}_v(E_v, \frac{1}{l})$  for all  $l \geq 1$  and  $v \in \mathfrak{M}_{\mathbb{K}}$ . Then, for all  $v \in \mathfrak{M}_{\mathbb{K}}$ , the sequence  $(\mu_{p_l, v})_{l \geq 1}$  converges to the equilibrium measure of  $\bar{E}_v$ .*

*Proof.* Our hypotheses imply that for every  $l \geq 1$  the Weil height of  $p_l$  is bounded from above independently of  $l$ . Together with the Northcott property and the fact that the points in the sequence  $(p_l)_{l \geq 1}$  are pairwise distinct, this implies that  $\lim_l \# \text{Gal}(p_l) = \infty$ . Taking a subsequence if necessary, we assume that  $\# \text{Gal}(p_l) \geq 2$  for every  $l \geq 1$ .

Since for each place  $v$  the space of Borel probability measures on  $\bar{\mathcal{B}}_v(E_v, 1)$  is sequentially compact, by taking a subsequence we can suppose without loss of generality that the sequence  $(\mu_{p_l, v})_{l \geq 1}$  converges to a probability measure  $\mu_v$  supported on  $\bigcap_l \bar{\mathcal{B}}_v(E_v, \frac{1}{l}) = \bar{E}_v$ .

For each  $l \geq 1$  and  $v \in \mathfrak{M}_{\mathbb{K}}$ , put for short  $G_{l, v} = \text{Gal}(p_l)_v$  and set

$$d_{l, v} = \frac{1}{\#G_{l, v}(\#G_{l, v} - 1)} \sum_{\substack{q, q' \in G_{l, v} \\ q \neq q'}} \log |q - q'|_v.$$

Consider also the probability measure on  $\mathbb{A}_v^{1, \text{an}} \times \mathbb{A}_v^{1, \text{an}}$ , given by

$$\nu_{l, v} = \frac{1}{\#G_{l, v}(\#G_{l, v} - 1)} \sum_{\substack{q, q' \in G_{l, v} \\ q \neq q'}} \delta_q \times \delta_{q'},$$

and note that  $(\nu_{l,v})_{l \geq 1}$  converges to  $\mu_v \times \mu_v$ . The function  $\log(\delta_v(\cdot, \cdot))$  is bounded from above on  $\mathcal{B}_v(E_v, 1) \times \mathcal{B}_v(E_v, 1)$ . Similarly as in the proof of Lemma 3.8, this property implies that

$$\begin{aligned} \limsup_{l \rightarrow \infty} d_{l,v} &= \limsup_{l \rightarrow \infty} \int_{\mathbb{A}_v^{1,\text{an}} \times \mathbb{A}_v^{1,\text{an}}} \log(\delta_v(z, z')) \, d\nu_{l,v}(z, z') \\ &\leq -I_v(\mu_v) \leq \log \text{cap}_v(\overline{E_v}). \end{aligned} \quad (7.4)$$

By the product formula,  $\sum_{v \in \mathfrak{M}_{\mathbb{K}}} n_v d_{l,v} = 0$ . Let  $S \subset \mathfrak{M}_{\mathbb{K}}$  be a finite set of places containing the Archimedean places and those where  $E_v \neq O_v$ . In particular,  $d_{l,v} \leq 0$  for  $v \notin S$ . Hence, for  $v \in \mathfrak{M}_{\mathbb{K}}$ ,

$$\begin{aligned} \liminf_{l \rightarrow \infty} d_{l,v} &= \liminf_{l \rightarrow \infty} \sum_{w \in \mathfrak{M}_{\mathbb{K}} \setminus \{v\}} -\frac{n_w}{n_v} d_{l,w} \\ &\geq \liminf_{l \rightarrow \infty} \sum_{w \in S \setminus \{v\}} -\frac{n_w}{n_v} d_{l,w} \\ &\geq - \sum_{w \in S \setminus \{v\}} \frac{n_w}{n_v} \limsup_{l \rightarrow \infty} d_{l,w} \\ &\geq - \sum_{w \in S \setminus \{v\}} \frac{n_w}{n_v} \log(\text{cap}_w(E_w)) \\ &\geq \log(\text{cap}_v(\overline{E_v})). \end{aligned}$$

Together with (7.4) and our hypothesis  $\text{cap}_v(\overline{E_v}) = \text{cap}_v(E_v)$ , this implies  $I_v(\mu_v) = -\log \text{cap}_v(\overline{E_v})$ . Therefore  $\mu_v$  is the equilibrium measure of  $\overline{E_v}$ , and the proof is complete.  $\square$

*Proof of Theorem 7.2.* Let  $(p_l)_{l \geq 1}$  be a sequence of pairwise distinct points of  $\overline{\mathbb{K}}^\times$  as in Proposition 7.4, which exists thanks to Proposition 7.3. Note in particular that the sequence  $(p_l)_{l \geq 1}$  is generic. On the other hand, Proposition 7.4 implies that, for every  $v \in \mathfrak{M}_{\mathbb{K}}$ , the sequence of probability measures  $(\mu_{p_l,v})_{l \geq 1}$  converges to  $\rho_{\overline{E_v}}$ . Here we have to show that, under the present hypotheses, this sequence of points is  $\overline{D}$ -small.

Let  $s_D$  be the canonical section of  $\mathcal{O}(D)$  with  $\text{div}(s_D) = D$ . This is a global section vanishing only at infinity. Hence for every  $v \in \mathfrak{M}_{\mathbb{K}}$  the  $v$ -adic Green function

$$g_{\overline{D},v} = -\log \|s_D\|_v$$

is a continuous real-valued function on  $\mathbb{A}_v^{1,\text{an}}$ . Let  $S \subset \mathfrak{M}_{\mathbb{K}}$  be a finite set of places containing the Archimedean places, the places where the metric  $\|\cdot\|_v$  differs from the canonical one, and those where  $E_v \neq O_v$ .

By construction, for each  $v \in \mathfrak{M}_{\mathbb{K}}$  and  $l \geq 1$  we have  $\text{Gal}(p_l)_v \subset \mathcal{B}_v(E_v, 1)$ . In particular, for  $v \notin S$ ,  $\text{Gal}(p_l)_v \subset O_v$  and so  $g_{\overline{D},v}(q) = 0$  for all  $q \in \text{Gal}(p_l)_v$ . Hence

$$h_{\overline{D}}(p_l) = \sum_{v \in \mathfrak{M}_{\mathbb{K}}} \frac{n_v}{\#\text{Gal}(p_l)_v} \sum_{q \in \text{Gal}(p_l)_v} g_{\overline{D},v}(q) = \sum_{v \in S} n_v \int \tilde{g}_{\overline{D},v} \, d\mu_{p_l,v}$$

for any continuous function  $\tilde{g}_{\overline{D},v}$  on  $\mathbb{P}_v^{1,\text{an}}$  coinciding with  $g_{\overline{D},v}$  on the bounded subset  $\mathcal{B}_v(E_v, 1)$ .

The measures  $\mu_{p_l,v}$  converge to  $\rho_{E_v}$  and are supported on the closure  $\overline{\mathcal{B}_v(E_v, 1)}$ . Also, for all  $v \notin S$ , we have  $\rho_{E_v} = \lambda_{\mathbb{S}_v,0}$  and  $g_{\overline{D},v}$  vanishes on the support of this measure. Hence

$$\lim_{l \rightarrow \infty} h_{\overline{D}}(p_l) = \sum_{v \in S} n_v \int \tilde{g}_{\overline{D},v} \, d\rho_{E_v} = \sum_{v \in \mathfrak{M}_{\mathbb{K}}} n_v \int g_{\overline{D},v} \, d\rho_{E_v}. \quad (7.5)$$

By the condition (3) and the fact that  $\mathbf{E}$  is an adelic set, we deduce that the collection  $\boldsymbol{\nu} = ((\text{val}_v)_* \rho_{E_v})_{v \in \mathfrak{M}_{\mathbb{K}}}$  is a centered adelic measure (Definition 4.4). Moreover,  $g_{\overline{D},v} = -\psi_{\overline{D},v} \circ \text{val}_v$  on  $\mathbb{A}_v^{1,\text{an}} \setminus \{0\}$ . By (7.5), we have

$$\lim_{l \rightarrow \infty} h_{\overline{D}}(p_l) = - \sum_{v \in \mathfrak{M}_{\mathbb{K}}} n_v \int \psi_{\overline{D},v} d(\text{val}_v)_* \rho_{E_v} = \eta_{\overline{D}}(\boldsymbol{\nu}).$$

Lemma 4.8 together with the conditions (1) and (2) implies that  $\eta_{\overline{D}}(\boldsymbol{\nu}) = \mu_{\overline{D}}^{\text{ess}}(X)$ . Hence the sequence  $(p_l)_{l \geq 1}$  is  $\overline{D}$ -small, as stated, finishing the proof of the theorem.  $\square$

**7.2. Local modulus concentration and equidistribution.** Corollary 4.13 gives a criterion for a semipositive toric metrized  $\mathbb{R}$ -divisor to satisfy the modulus concentration property at a given place. Applying it, one can immediately give examples where modulus concentration fails at that place. If this happens, then the equidistribution property also fails at that place.

Can this absence of modulus concentration affect the equidistribution property at another place? The next result on the projective line over a number field shows that this can be the case under a rationality hypothesis.

**Proposition 7.5.** *Let  $X = \mathbb{P}_{\mathbb{K}}^1$  be the projective line over a number field  $\mathbb{K}$ ,  $\overline{D}$  the divisor at infinity equipped with a semipositive toric metric, and  $v_0 \in \mathfrak{M}_{\mathbb{K}}$ . For each  $v \in \mathfrak{M}_{\mathbb{K}}$ , let  $B_v$  be the set introduced in Notation 4.2. Assume that there is a point  $p \in X_0(\overline{\mathbb{K}}) = \overline{\mathbb{K}}^{\times}$  such that  $\text{val}_v(p) \in B_v$  for all  $v \in \mathfrak{M}_{\mathbb{K}}$  and  $\text{val}_{v_0}(p) \in \text{ri}(B_{v_0})$ .*

*If  $\overline{D}$  does not satisfy the modulus concentration property at  $v_0$ , then  $\overline{D}$  does not satisfy the equidistribution property at any place of  $\mathbb{K}$ .*

*Proof.* Assume that  $\overline{D}$  does not satisfy the modulus concentration property at  $v_0$ . Let  $v \in \mathfrak{M}_{\mathbb{K}}$ . If  $v = v_0$  then clearly  $\overline{D}$  does not satisfy the equidistribution property at  $v$ , so we can suppose that  $v \neq v_0$ . Extending scalars to a suitable large number field and translating by the point  $p$ , we can also reduce to the case when  $0 \in \text{ri}(B_{v_0})$  and  $0 \in B_w$  for all  $w \in \mathfrak{M}_{\mathbb{K}}$ .

Let  $F_{v_0}$ ,  $g_{1,v_0}$  and  $g_{2,v_0}$  be as in Notation 4.2, and let  $x$  be a in  $\Delta_D$  at which  $g_{1,v_0} + g_{2,v_0}$  attains its maximum. By Corollary 4.13, the set  $F_{v_0}$  is not a single point. Since  $0 \in \text{ri}(B_{v_0})$  and  $F_{v_0}$  is the minimal face of  $\partial g_{1,v_0}(x)$  containing  $B_{v_0}$ , there is  $\delta > 0$  such that the set  $F_{v_0}$  contains the interval  $[-\delta, \delta]$ . Set

$$c = \frac{e^{\delta} + e^{-\delta}}{2} > 1$$

and consider the closed bounded adelic set  $\mathbf{E} = (E_w)_{w \in \mathfrak{M}_{\mathbb{K}}}$  given by

$$E_{v_0} = \begin{cases} [-2c, 2c] & \text{if } v_0 \text{ is Archimedean,} \\ \mathcal{B}_{v_0}(2, c) & \text{if } v_0 \text{ is non-Archimedean,} \end{cases}$$

$$E_v = \begin{cases} [-2/c, 2/c] & \text{if } v \text{ is Archimedean,} \\ \mathcal{B}_v(2, 1/c) & \text{if } v \text{ is non-Archimedean,} \end{cases}$$

and, for  $w \neq v_0, v$ ,

$$E_w = \begin{cases} [-2, 2] & \text{if } w \text{ is Archimedean,} \\ \mathcal{O}_w = \mathcal{B}_w(0, 1) & \text{if } w \text{ is non-Archimedean.} \end{cases}$$

The local capacities of these sets are

$$\text{cap}_{v_0}(E_{v_0}) = c, \quad \text{cap}_v(E_v) = 1/c \quad \text{and} \quad \text{cap}_w(E_w) = 1 \quad \text{for } w \neq v_0, v,$$

see for instance [Rum02, §3]. Hence, the global capacity of  $\mathbf{E}$  is 1.

Consider the map  $R: \mathbb{P}_{\mathbb{K}}^1 \rightarrow \mathbb{P}_{\mathbb{K}}^1$  defined in affine coordinates by  $R(z) = z + \frac{1}{z}$ . Using the expression  $R(z) - 2 = \frac{(z-1)^2}{z}$ , one checks that, for  $w$  non-Archimedean,

$$\begin{aligned} R^{-1}(E_w) &= \begin{cases} \{z \in \mathbb{C}_{v_0} \mid |z-1|_{v_0}^2 \leq c|z|_{v_0}\} & \text{if } w = v_0, \\ \{z \in \mathbb{C}_v \mid |z-1|_v^2 \leq c^{-1}|z|_v\} & \text{if } w = v, \\ \{z \in \mathbb{C}_w \mid |z^2+1|_w \leq |z|_w\} & \text{if } w \neq v_0, v, \end{cases} \\ &= \begin{cases} \{z \in \mathbb{C}_{v_0} \mid c^{-1} \leq |z|_{v_0} \leq c\} & \text{if } w = v_0, \\ \mathcal{B}_v(1, c^{-1/2}) & \text{if } w = v, \\ \mathbb{S}_w & \text{if } w \neq v_0, v. \end{cases} \end{aligned} \quad (7.6)$$

On the other hand, using

$$z = \frac{1}{2} \left( R(z) \pm \sqrt{R(z)^2 - 4} \right), \quad c - \sqrt{c^2 - 1} = e^{-\delta} \quad \text{and} \quad c + \sqrt{c^2 - 1} = e^{\delta},$$

one also checks that, for  $w$  Archimedean,

$$R^{-1}(E_w) = \begin{cases} \mathbb{S}_{v_0} \cup \{z \in \mathbb{C}_{v_0} \mid \text{im}(z) = 0, e^{-\delta} \leq |z|_{v_0} \leq e^{\delta}\} & \text{if } w = v_0, \\ \{z \in \mathbb{S}_v \mid \text{im}(z) \geq \sqrt{1-c^{-2}}\} & \text{if } w = v, \\ \mathbb{S}_w & \text{if } w \neq v_0, v. \end{cases} \quad (7.7)$$

We represent in Figure 4 the inverse images by  $R$  of the sets  $E_{v_0}$ ,  $E_v$  and  $E_w$  in the Archimedean case. The point  $x$  therein is  $x = c^{-1} + i\sqrt{1-c^{-2}}$ .




FIGURE 4. Inverse images by  $R$  of the sets  $E_{v_0}$ ,  $E_v$  and  $E_w$  for  $v_0$ ,  $v$  and  $w \neq v, v_0$  Archimedean

We deduce from the previous analysis that, regardless whether  $v_0$ ,  $v$  or  $w$  are Archimedean or not, we have

$$R^{-1}(E_{v_0}) \subset \text{val}_{v_0}^{-1}([-\delta, \delta]), \quad R^{-1}(E_v) \subsetneq \mathbb{S}_v \quad \text{and} \quad R^{-1}(E_w) = \mathbb{S}_w \text{ for } w \neq v_0, v.$$

Let  $(p_l)_{l \geq 1}$  be a sequence of pairwise distinct points as given by Proposition 7.3 applied to the adelic set  $\mathbf{E}$ . For each  $l \geq 1$ , choose a point  $q_l \in R^{-1}(p_l)$ . Since for each place  $v$  the space of Borel probability measures on  $\overline{\mathcal{B}}_v(E_v, 1)$  is sequentially compact, after restricting to a subsequence we can assume that the sequence  $(\mu_{q_l, w})_{l \geq 1}$  converges to a probability measure  $\mu_w$  on  $\overline{R^{-1}(E_w)}$ , for all  $w \in \mathfrak{M}_{\mathbb{K}}$ . By construction, for each  $w$  the supports of the direct image measures  $\nu_{q_l, w} = (\text{val}_w)_* \mu_{q_l, w}$ ,  $l \geq 1$ , are contained in  $[-\delta, \delta] \subset N_{\mathbb{R}}$ . Therefore, this sequence of measures converges in the KR-topology to the direct image  $(\text{val}_w)_* \mu_w$ , which can be seen by using Remark 3.13.

Let  $S \subset \mathfrak{M}_{\mathbb{K}}$  be the finite subset consisting of the Archimedean places plus  $v_0$  and  $v$ . If  $w \neq v_0$ , then  $\text{Gal}(q_l)_w \subset \text{val}_w^{-1}(0)$  and  $E[\nu_{q_l, w}] = 0$ . Thus

$$E[(\text{val}_w)_*(\mu_w)] = \lim_l E[\nu_{q_l, w}] = 0.$$

Hence, thanks to the convergence in the KR-topology and the product formula,

$$E[(\text{val}_{v_0})_*(\mu_{v_0})] = \lim_l E[\nu_{q_l, v_0}] = \lim_l \sum_{\substack{w \in S \\ w \neq v_0}} -E[\nu_{q_l, w}] = 0.$$

Thus  $E[(\text{val}_w)_*(\mu_w)] = 0 \in B_w$  for all  $w \in \mathfrak{M}_{\mathbb{K}}$ . By construction, it is also clear that  $\text{supp}((\text{val}_w)_*\mu_w) \subset F_w$  for all  $w$ . By Lemma 4.8, the sequence  $(q_l)_{l \geq 1}$  is  $\overline{D}$ -small.

We have thus constructed a generic  $\overline{D}$ -small sequence such that its  $v$ -adic Galois orbit converges to a measure  $\mu_v$  whose support is contained in the closure  $\overline{R^{-1}(E_v)}$ . On the other hand, the sequence  $(\omega_l)_{l \geq 1}$  given by the choice of a primitive  $l$ -th root of unity is also  $\overline{D}$ -small, but its  $v$ -adic Galois orbit converges to the measure  $\lambda_{\mathbb{S}, 0}$ . By (7.6) and (7.7) the support of this measure is not contained in  $\overline{R^{-1}(E_v)}$ , so it is different from  $\mu_v$ . We deduce that  $\overline{D}$  does not satisfy the  $v$ -adic equidistribution property, as stated.  $\square$

**Example 7.6.** Let  $X = \mathbb{P}_{\mathbb{Q}}^1$  and  $\overline{D}$  the divisor at infinity plus the divisor at zero, equipped with the semipositive toric metric from Example 6.2. As explained therein,  $\overline{D}$  does not satisfy modulus concentration at the place  $v_0 = 2$  and, by (6.1), we have  $0 \in \text{ri}(B_v)$  for all  $v \in \mathfrak{M}_{\mathbb{Q}}$ . Theorem 7.2 implies that  $\overline{D}$  does not satisfy the equidistribution property for any place of  $\mathbb{Q}$ .

**Remark 7.7.** A rationality hypothesis like the condition that the sets  $B_v$  contain the image by the valuations map of an algebraic point, is necessary for the conclusion of Proposition 7.5 to hold. Indeed, suppose that, for a given non-Archimedean place  $v$ , we have  $B_v = F_v = \{u_v\}$  with  $u_v \notin \text{val}_v(\overline{\mathbb{K}}_v^{\times})$ . By the tree structure of the Berkovich projective line, this implies that  $\text{val}_v^{-1}(u_v)$  consists of a single point, of type III in Berkovich's classification [BR10, §1.4]. Hence, the  $v$ -adic modulus concentration at  $v$  given by Corollary 4.13, easily implies that the  $v$ -adic Galois orbits of  $\overline{D}$ -small sequences of algebraic points concentrate around this point of type III, regardless of the structure of the set  $B_{v_0}$ .

## REFERENCES

- [AW45] E. Artin and G. Whaples, *Axiomatic characterization of fields by the product formula for valuations*, Bull. Amer. Math. Soc. **51** (1945), 469–492.
- [BB10] R. Berman and S. Boucksom, *Growth of balls of holomorphic sections and energy at equilibrium*, Invent. Math. **181** (2010), 337–394.
- [Ber90] V. G. Berkovich, *Spectral theory and analytic geometry over non-Archimedean fields*, Math. Surveys Monogr., vol. 33, Amer. Math. Soc., 1990.
- [Bil97] Y. Bilu, *Limit distribution of small points on algebraic tori*, Duke Math. J. **89** (1997), 465–476.
- [BMPS16] J. I. Burgos Gil, A. Moriwaki, P. Philippon, and M. Sombra, *Arithmetic positivity on toric varieties*, J. Alg. Geom. **25** (2016) 201–272.

- [Bou85] N. Bourbaki, *Éléments de mathématique. Fasc. XXX. Algèbre commutative. Chapitre 5: Entiers. Chapitre 6: Valuations*, Actualités Scientifiques et Industrielles, No. 1308 Hermann, 1964.
- [Bou69] ———, *Éléments de mathématique. Fasc. XXXV. Livre VI: Intégration. Chapitre IX: Intégration sur les espaces topologiques séparés*, Actualités Scientifiques et Industrielles, No. 1343, Hermann, 1969.
- [BPS14] J. I. Burgos Gil, P. Philippon, and M. Sombra, *Arithmetic geometry of toric varieties. Metrics, measures and heights*, Astérisque, vol. 360, Soc. Math. France, 2014.
- [BPS15] ———, *Successive minima of toric height functions*, Ann. Inst. Fourier (Grenoble) **65** (2015), 2145–2197.
- [BR06] M. Baker and R. Rumely, *Equidistribution of small points, rational dynamics, and potential theory*, Ann. Inst. Fourier (Grenoble) **56** (2006), 625–688.
- [BR10] ———, *Potential theory and dynamics on the Berkovich projective line*, Math. Surveys Monogr., vol. 159, Amer. Math. Soc., 2010.
- [Cha00] A. Chambert-Loir, *Points de petite hauteur sur les variétés semi-abéliennes*, Ann. Sci. Éc. Norm. Sup. **33** (2000), 789–821.
- [Cha06] ———, *Mesures et équidistribution sur les espaces de Berkovich*, J. Reine Angew. Math. **595** (2006), 215–235.
- [Che11] H. Chen, *Differentiability of the arithmetic volume function*, J. Lond. Math. Soc. (2) **84** (2011), 365–384.
- [Cin11] Z. Cinkir, *Zhang’s conjecture and the effective Bogomolov conjecture over function fields*, Invent. Math. **183** (2011), 517–562.
- [Fab09] X. Faber, *Equidistribution of dynamically small subvarieties over the function field of a curve*, Acta Arith. **137** (2009), 345–389.
- [FR06] C. Favre and J. Rivera-Letelier, *Équidistribution quantitative des points de petite hauteur sur la droite projective*, Math. Ann. **335** (2006), 311–361, *corrigendum* in Math. Ann. **339** (2007), 799–801.
- [Ghi09] D. Ghioca, *Points of small height on varieties defined over a function field*, Canad. Math. Bull. **52** (2009), 237–244.
- [Gub07] W. Gubler, *The Bogomolov conjecture for totally degenerate abelian varieties*, Invent. Math. **169** (2007), 377–400.
- [Gub08] ———, *Equidistribution over function fields*, Manuscripta Math. **127** (2008), 485–510.
- [Liu02] Q. Liu, *Algebraic geometry and arithmetic curves*, Oxf. Grad. Texts Math., vol. 6, Oxford Univ. Press, 2002.
- [Roc70] R. T. Rockafellar, *Convex analysis*, Princeton Math. Series, vol. 28, Princeton Univ. Press, 1970.
- [Rum02] R. Rumely, *The Fekete-Szegő theorem with splitting conditions. II*, Acta Arith. **103** (2002), 347–410.
- [SUZ97] L. Szpiro, E. Ullmo, and S.-W. Zhang, *Équirépartition des petits points*, Invent. Math. **127** (1997), 337–347.
- [Tsu75] M. Tsuji, *Potential theory in modern function theory*, Chelsea Publishing Co., 1975.
- [Ull98] E. Ullmo, *Positivité et discrétion des points algébriques des courbes*, Annals of Math. **147** (1998), 167–179.
- [Vil09] C. Villani, *Optimal transport. Old and new*, Grundlehren Math. Wiss., vol. 338, Springer-Verlag, 2009.
- [Wei74] A. Weil, *Basic number theory*, third ed., Grundlehren Math. Wiss., vol. 144, Springer-Verlag, 1974.
- [Yam13] K. Yamaki, *Geometric Bogomolov conjecture for abelian varieties and some results for those with some degeneration (with an appendix by Walter Gubler: the minimal dimension of a canonical measure)*, Manuscripta Math. **142** (2013), 273–306.
- [Yam16] K. Yamaki, *Strict supports of canonical measures and applications to the geometric Bogomolov conjecture*, Compos. Math. **152** (2016), 997–1040.
- [Yua08] X. Yuan, *Big line bundles over arithmetic varieties*, Invent. Math. **173** (2008), 603–649.
- [Zha95] S.-W. Zhang, *Positive line bundles on arithmetic varieties*, J. Amer. Math. Soc. **8** (1995), 187–221.
- [Zha98] ———, *Equidistribution of small points of abelian varieties*, Annals of Math. **147** (1998), 159–165.

INSTITUTO DE CIENCIAS MATEMÁTICAS (CSIC-UAM-UCM-UCM3). CALLE NICOLÁS CABRE-  
RA 15, CAMPUS UAM, CANTOBLANCO, 28049 MADRID, SPAIN

*E-mail address:* [burgos@icmat.es](mailto:burgos@icmat.es)

*URL:* <http://www.icmat.es/miembros/burgos>

INSTITUT DE MATHÉMATIQUES DE JUSSIEU – U.M.R. 7586 DU CNRS, ÉQUIPE DE THÉORIE DES  
NOMBRES. BP 247, 4 PLACE JUSSIEU, 75005 PARIS, FRANCE

*E-mail address:* [patrice.philippon@imj-prg.fr](mailto:patrice.philippon@imj-prg.fr)

*URL:* <http://www.math.jussieu.fr/~pph>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER. HYLAN BUILDING, ROCHESTER,  
NY 14627, U.S.A.

*E-mail address:* [riversletelier@gmail.com](mailto:riversletelier@gmail.com)

*URL:* <http://rivers-letelier.org>

INSTITUCIÓ CATALANA DE RECERCA I ESTUDIS AVANATS (ICREA). PASSEIG LLUÍS COMPA-  
NYS 23, 08010 BARCELONA, SPAIN

DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, UNIVERSITAT DE BARCELONA (UB). GRAN  
VIA 585, 08007 BARCELONA, SPAIN

*E-mail address:* [sombra@ub.edu](mailto:sombra@ub.edu)

*URL:* <http://www.maia.ub.edu/~sombra>